

LARGE CARDINALS AND DOWKER PRODUCTS

CHRIS GOOD

May 1993

ABSTRACT. We prove that if there is a model of set-theory which contains no first countable, locally compact, scattered, countably paracompact space X , whose Tychonoff square is a Dowker space, then there is an inner model which contains a measurable cardinal.

In this paper we always take space to mean Hausdorff topological space. A space is normal if every pair of disjoint closed sets can be separated by disjoint open sets, and binormal if its product with the closed unit interval is normal. A space is countably paracompact (metacompact) if every countable open cover has a locally (point) finite open refinement. In [Dk], Dowker shows that a normal space is binormal iff it is countably paracompact iff it is countably metacompact. A Dowker space is a normal space that is not countably paracompact. For a survey of Dowker spaces we refer the reader to [R].

Rudin and Starbird [RS] have shown that for normal, countably paracompact X and metrizable M , $X \times M$ is normal if and only if it is countably paracompact. They asked whether a product of two normal, countably paracompact spaces could be a Dowker space. Bešlagić constructs various positive answers to this question, assuming $\diamond$ or CH , in [B1,B2&B3].

In [G] we prove that if there is a model of set theory which contains no first countable, locally compact, scattered Dowker spaces, then there is a model of set-theory which contains a measurable cardinal. Here we extend this result by proving that large cardinals are needed for a model in which there is no first countable, locally compact, countably paracompact space X with first countable, locally compact, scattered Dowker square:

1. Theorem. *If no inner model of set theory contains a measurable cardinal, then there is a first countable, locally countable, locally compact, strongly zero-dimensional, collectionwise normal, countably paracompact, scattered space whose Tychonoff square is a first countable, locally compact, collectionwise normal, scattered Dowker space.*

Notation and terminology are standard—see [E], [K] or [KV]. We regard cardinals as initial ordinals, and an ordinal as the set of its predecessors. We use the term *club set* or *club* to denote a closed, unbounded subset of an ordinal—it will be clear from the context which particular ordinal we mean. For a function $f : A \rightarrow B$, we denote by $f \restriction C$ the set $\{f(x) : x \in C \subseteq A\}$. For a subset A of $\alpha \times \beta$, we denote the set $\{\gamma : (\exists \delta)(\gamma, \delta) \in A\}$ by $\text{dom } A$, and the set $\{\delta : (\exists \gamma)(\gamma, \delta) \in A\}$ by $\text{ran } A$. Following [B1], we call a subset A of $\kappa^+ \times \kappa^+$ is said to be 2-unbounded if A is not a subset of $(\kappa^+ \times \alpha) \cup (\alpha \times \kappa^+)$ for any $\alpha \in \kappa^+$. As usual we use the following characterization from [Dk]:

1991 *Mathematics Subject Classification.* 03E35, 03E55, 54D15, 54D20, 54EG15.

Key words and phrases. small Dowker space, Dowker product, normality, countable paracompactness, measurable cardinal, Covering Lemma.

a space is countably metacompact if and only if, for every decreasing sequence $\{D_n\}_{n \in \omega}$ of closed subsets of X , which has empty intersection, there is a sequence $\{U_n\}_{n \in \omega}$ of open sets, U_n containing D_n for each n , which also has empty intersection.

A stationary subset E of some uncountable cardinal λ is said to be *non-reflecting* if, for every $\alpha < \lambda$, $\alpha \cap E$ is non-stationary in α . If E is a non-reflecting stationary subset of κ^+ and $\alpha \in \kappa^+$, then it is easy to see that there is a club set $H = \{\gamma_\lambda : \lambda \in \theta \leq \alpha\}$ of α such that H and E are disjoint, and $(\gamma_\lambda, \gamma_{\lambda+1})$ is countable for all $\lambda \in \theta$. In what follows we shall let E denote a non-reflecting, stationary subset of κ^+ , each member of which has countable cofinality

2. Definition. $\clubsuit_{\kappa^+}(E, 2)$ is the assertion that there is a collection $\{R_{\alpha,i} : R_{\alpha,i} \subseteq \alpha, \alpha \in E \cap \text{LIM and } i \in 2\}$ such that each $R_{\alpha,i}$ is an ω -sequence, cofinal in α , and $\{\alpha \in E : R_{\alpha,i} \subseteq X_i \text{ for both } i \in 2\}$ is stationary whenever X_0 and X_1 are unbounded subsets of κ^+ .

In [G] we deduce, via [Dv], [DJ] and [F],

3. Lemma. *If no inner model of set-theory contains a measurable cardinal, then $\clubsuit_{\kappa^+}(E, 2)$ for some κ^+ . $\square$*

In the construction of the space X , we use the following two consequences of $\clubsuit_{\kappa^+}(E, 2)$.

4. Definition. $\clubsuit_{\kappa^+ \times \kappa^+}(E, 2)$ is the assertion that there is a sequence $\{S_{\alpha,i} : S_{\alpha,i} \subseteq \alpha \times \alpha, \alpha \in E \cap \text{LIM and } i \in 2\}$ such that $S_{\alpha,i}$ is an ω -sequence, cofinal in $\alpha \times \alpha$, and $\{\alpha \in E : S_{\alpha,i} \subseteq X_i \text{ for both } i \in 2\}$ is stationary whenever X_0 and X_1 are 2-unbounded subsets of $\kappa^+ \times \kappa^+$.

5 Definition. $\clubsuit_{\kappa^+}^\cap(E, 2)$ is the assertion that there is a sequence $\{T_{\alpha,i,n} : T_{\alpha,i,n} \subseteq \alpha, \alpha \in E \cap \text{LIM, and } i \in 2\}$ such that $\bigcup_{n \in \omega} T_{\alpha,i,n}$ and each $T_{\alpha,i,n}$ is an ω -sequence, cofinal in α , $T_{\alpha,i,n} \cap T_{\alpha,j,m}$ is empty whenever $i \neq j$ or $m \neq n$, and $\bigcap_{n \in \omega} \{\alpha \in E : T_{\alpha,i,n} \subseteq X_{i,n}, \text{ for both } i \in 2\}$ is stationary whenever $\{X_{i,n} : i \in 2, n \in \omega\}$ is a collection of unbounded subsets of κ^+ .

6. Lemma. *If $\clubsuit_{\kappa^+}(E, 2)$, then $\clubsuit_{\kappa^+ \times \kappa^+}(E, 2)$ and $\clubsuit_{\kappa^+}^\cap(E, 2)$*

Proof. Let $\{R_{\alpha,i} : \alpha \in E \cap \text{LIM, } i \in 2\}$ be a $\clubsuit_{\kappa^+}(E, 2)$ -sequence. We may assume that $R_{\alpha,0}$ and $R_{\alpha,1}$ are disjoint for all α in $E \cap \text{LIM}$. Let $f : \kappa^+ \rightarrow \kappa^+ \times \kappa^+$ and $g : \kappa^+ \rightarrow \kappa^+ \times \omega$ be any bijections. $F = \{\alpha : f''\alpha = \alpha \times \alpha\}$ and $G = \{\alpha : g''\alpha = \alpha \times \omega\}$ are both club in κ^+ .

For α in $E \cap F \cap \text{LIM}$ such that both $f''R_{\alpha,0}$ and $f''R_{\alpha,1}$ are cofinal in $\alpha \times \alpha$, define $S_{\alpha,i}$ to be the set $f''R_{\alpha,i}$. Otherwise, for α in E , let $S_{\alpha,i}$ be an arbitrary sequence cofinal in $\alpha \times \alpha$. It is easy to see that $\{S_{\alpha,i} : S_{\alpha,i} \subseteq \alpha \times \alpha, \alpha \in E \cap \text{LIM and } i \in 2\}$ is a $\clubsuit_{\kappa^+ \times \kappa^+}(E, 2)$ -sequence.

If α is in $E \cap G \cap \text{LIM}$, $i \in 2$ and $n \in \omega$, let $T_{\alpha,i,n}$ be the set $\text{dom}(g''B_{\alpha,i} \cap (\alpha \times \{n\}))$. Otherwise, for α in E , let $T_{\alpha,i,n}$ be arbitrary.

To see that $\{T_{\alpha,i,n} : T_{\alpha,i,n} \subseteq \alpha, \alpha \in E \cap \text{LIM, and } i \in 2\}$ is a $\clubsuit_{\kappa^+}^\cap(E, 2)$ -sequence, let $\{X_{i,n}\}_{i \in 2, n \in \omega}$ be a collection of unbounded subsets of κ^+ , and let $X_i = \bigcup_{n \in \omega} X_{i,n} \times \{n\}$. $S = \{\alpha \in E : R_{\alpha,i} \subseteq g^{-1}''X_i, i \in 2\}$ is stationary. If α is in S then $g''R_{\alpha,i}$ is a subset of X_i and hence S is a subset of $\bigcap \{\alpha \in E : T_{\alpha,i,n} \subseteq X_{i,n}, i \in 2\}$. $\square$

Our construction is similar to that used by Bešlagić in [B1]. We define three normal topologies, $\mathcal{T}_i$, $i \in 3$, on the point set $Y = \kappa^+ \times \omega$. The topologies $\mathcal{T}_0$ and $\mathcal{T}_1$ both refine $\mathcal{T}_2$, which is a Hausdorff topology, hence the diagonal Δ of $(Y, \mathcal{T}_0) \times (Y, \mathcal{T}_1)$ is a closed subspace of X^2 . Our space X is the disjoint topological sum of $(Y, \mathcal{T}_0)$ and $(Y, \mathcal{T}_1)$. $\clubsuit_{\kappa^+ \times \kappa^+}(E, 2)$ helps to ensure that the product X^2 is normal, and that Δ is a Dowker space. Since Δ is closed in X^2 , X^2 is also a Dowker space. We use $\clubsuit_{\kappa^+}^\cap(E, 2)$ to ensure that $(Y, \mathcal{T}_i)$, $i \in 2$ is countably paracompact (cf §5 [B1]).

7. Example. $\clubsuit_{\kappa^+}(E, 2)$ *There is a first countable, locally countable, locally compact, strongly zero-dimensional, collectionwise normal, countably paracompact, scattered space X , whose Tychonoff square is a first countable locally compact, collectionwise normal, scattered Dowker space.*

Proof. Let Y be the point set $\kappa^+ \times \omega$, let $\pi : Y \rightarrow \kappa^+$ be the natural projection, $\pi((\alpha, n)) = \alpha$, and let $\Pi : Y^2 \rightarrow \kappa^{+2}$ be the natural projection, $\Pi((\alpha, n), (\beta, m)) = (\alpha, \beta)$. Let $\{S_{\alpha, i} : S_{\alpha, i} \subseteq \alpha \times \alpha, \alpha \in E \cap \text{LIM and } i \in 2\}$ and $\{T_{\alpha, i, n} : T_{\alpha, i, n} \subseteq \alpha, \alpha \in E \cap \text{LIM}, n \in \omega \text{ and } i \in 2\}$ be $\clubsuit_{\kappa^+ \times \kappa^+}(E, 2)$ - and $\clubsuit_{\kappa^+}(E, 2)$ -sequences respectively. Bearing in mind the proof of Lemma 6, it is not hard to see that we may assume that $\bigcup_{i \in 2} (\text{ran } S_{\alpha, i} \cup \text{dom } S_{\alpha, i})$ and $\bigcup_{n \in \omega} T_{\alpha, i, n}$ are disjoint for each α in $E \cap \text{LIM}$. We may also assume that each ω -sequence $S_{\alpha, i}$ is strictly increasing in both coordinates.

For each α in $E \cap \text{LIM}$ and each $i \in 2$, partition $S_{\alpha, i}$ into ω disjoint sequences $S_{\alpha, i, n}$, where $n \in \omega$, each cofinal in $\alpha \times \alpha$. Let $B(\alpha, i, n)$ be the sequence $\text{dom } S_{\alpha, i, n} \cup \text{ran } S_{\alpha, i, n}$. For each $n \in \omega$, $B(\alpha, i, n)$ is an ω -sequence, cofinal in α , and the collection $\{B(\alpha, i, n) : \alpha \in E \cap \text{LIM}, i \in 2\}$ is a $\clubsuit_{\kappa^+}(E, 2)$ -sequence. Since $S_{\alpha, i}$ is strictly increasing in both coordinates, $B(\alpha, i, n)$ and $B(\alpha, i, m)$ are disjoint whenever $n \neq m$. Let $B(\alpha, n) = B(\alpha, 0, n) \cup B(\alpha, 1, n)$ and let $B(\alpha) = \bigcup_{n \in \omega} B(\alpha, n)$. Enumerate the ω -sequence $B(\alpha)$ increasingly as $\{\beta(\alpha, j) : j \in \omega\}$.

For $i \in 2$ let $C(\alpha, i, n) = T_{\alpha, i, n}$ and let $C(\alpha, 2, n) = C(\alpha, 0, n) \cup C(\alpha, 1, n)$. Let $C(\alpha) = \bigcup_{n \in \omega} C(\alpha, 2, n)$. Enumerate the ω -sequence $C(\alpha)$ increasingly as $\{\gamma(\alpha, j) : j \in \omega\}$.

By assumption $B(\alpha)$ and $C(\alpha)$ are disjoint for all α in $E \cap \text{LIM}$. Let $A(\alpha) = B(\alpha) \cup C(\alpha)$ and index $A(\alpha)$ increasingly as $\{\alpha(k) : k \in \omega\}$.

We define the topologies $\mathcal{T}_i$ by induction on the lexicographical order on $\kappa^+ \times \omega$. At each stage of the induction (α, n) , and for each $i \in 3$, we define a topology $\mathcal{T}_{i, \alpha}$ on $Y_\alpha = \alpha \times \omega$ and then a neighbourhood base $\mathcal{N}_i(\alpha, n) = \{N_i((\alpha, n), k)\}_{k \in \omega}$ at the point (α, n) . Our inductive hypotheses are, for $\gamma < \beta < \alpha$ and $i \in 3$:

- (1) $\mathcal{T}_{i, \beta}$ is a Hausdorff, conservative extension of $\mathcal{T}_{i, \gamma}$, and $Y_{\gamma+1}$ is a $\mathcal{T}_{i, \beta}$ -clopen subset of Y_β ;
- (2) $\mathcal{N}_i(\gamma, k)$ is a decreasing neighbourhood base of sets which are clopen, compact and countable under $\mathcal{T}_{i, \beta}$, and are subsets of $Y_{\gamma+1}$;
- (3) $N_i((\beta, n), k)$ and $N_i((\beta, m), k)$ are disjoint whenever $n \neq m$;
- (4) $N_0((\beta, n), k) \cup N_1((\beta, n), k)$ is a subset of $N_2((\beta, n), k)$ for all $k \in \omega$;
- (5) if $\delta(n, k) = \inf\{\pi"N_i((\alpha, n), k)\}$, then, for all $n \in \omega$, the sequence $\{\delta(n, k) : k \in \omega\}$ is cofinal in α ;
- (6) for all $0 < r \in \omega$, the point $(\gamma, 0)$ is a $\mathcal{T}_{0, \beta}$ -limit of each sequence $C(\gamma, 0, r) \times \{r\}$, a $\mathcal{T}_{1, \beta}$ -limit of each sequence $C(\gamma, 1, r) \times \{r\}$, and a $\mathcal{T}_{2, \beta}$ -limit of both sequences;
- (7) if $N_0 \in \mathcal{N}_0(\beta, 0)$ and $N_1 \in \mathcal{N}_1(\beta, 0)$, then $N_0 \cap N_1 = \{(\beta, 0)\}$;
- (8) for all $0 \leq p \leq m$, the point $(\beta, m+1)$ is a $\mathcal{T}_i$ -limit of the sequence $B(\alpha, m) \times \{p\}$.

If $\alpha = 0$, let $\mathcal{T}_{i, 0} = \emptyset$ and let $\mathcal{N}_i(0, n) = \{\{(0, n)\}\}$ for each $i \in 3$. Suppose that we have defined $\mathcal{N}_i(\beta, k)$ for each $i \in 3$, all $\beta \in \alpha$ and all $k \in \omega$. Define $\mathcal{T}_{i, \alpha}$ to be the topology generated by $\bigcup\{\mathcal{N}_i(\beta, k) : k \in \omega, \beta < \alpha\}$.

If $\alpha = \beta + 1$ for some β , or α is not in E , then we declare the point (α, n) to be isolated and define $\mathcal{N}_i(\alpha, n)$ to be $\{\{(\alpha, n)\}\}$ for each $i \in 3$.

Now suppose that α is a limit ordinal in E .

First let us suppose that $n = 0$. The sequence $C(\alpha)$ is enumerated as $\{\gamma(\alpha, j) : j \in \omega\}$. Each $\gamma(\alpha, j)$ in $C(\alpha)$ occurs uniquely in T_{α, i_j, r_j} for some $i_j \in 2$ and some $r_j \in \omega$, and is indexed as $\alpha(k_j)$ in $A(\alpha)$. By inductive hypotheses (4) and (5), whenever $r_j > 0$, we can choose a basic open set $N_2(\gamma(\alpha, j), r_j)$ from $\mathcal{N}_2(\gamma(\alpha, j), r_j)$ such that

- (†) $\pi"N_2(\gamma(\alpha, j), r_j)$ is a subset of the interval $(\alpha(k_j - 1), \alpha(k_j)]$ in κ^+ (by (5)).

For $i \in 3$, and each $k \in \omega$, define

$$N_i((\alpha, 0), k) = \{(\alpha, 0)\} \cup \bigcup\{N_2(\gamma(\alpha, j), r_j) : \gamma(\alpha, j) \in C(\alpha, i, r_j), r_j > 0, j > k\}.$$

Now suppose that $n = m + 1$ for some $m \in \omega$. The sequence $B(\alpha)$ is enumerated as $\{\beta(\alpha, j) : j \in \omega\}$, and each $\beta(\alpha, j)$ occurs uniquely in some $B(\alpha, r_j)$, and is indexed in $A(\alpha)$ as $\alpha(k_j)$. By (4),

(5) and the fact that $\mathcal{T}_{2\alpha}$ is Hausdorff, for each $\beta(\alpha, r_j)$ such that $r_j = m$ and for each $p \leq r_j$, we can choose disjoint basic open neighbourhoods $N_i(\beta(\alpha, j), p)$ from $\mathcal{N}_i(\beta(\alpha, j), p)$, $i \in 3$ of the point $(\beta(\alpha, j), p)$ such that $N_i(\beta(\alpha, j), p)$ is a subset of $N_2(\beta(\alpha, j), p)$, for each $i \in 2$, and

($\ddagger$) $\pi"N_i(\beta(\alpha, j), p)$ is a subset of the interval $(\alpha(k_j - 1), \alpha(k_j)]$ in κ^+ .

For $i \in 3$, and each $k \in \omega$, define

$$N_i((\alpha, n), k) = \{(\alpha, n)\} \cup \bigcup \{N_i(\beta(\alpha, j), p) : \beta(\alpha, j) \in B(\alpha, m), p \leq m, \text{ and } j > k\}.$$

It is not hard to check that the inductive hypotheses hold

Let $\mathcal{T}_i$ be the topology generated by $\bigcup_{(\alpha, n) \in Y} \mathcal{N}_i(\alpha, n)$.

Clearly both $\mathcal{T}_0$ and $\mathcal{T}_1$ refine $\mathcal{T}_2$, and it is not hard to check that each $(Y, \mathcal{T}_i)$ is Hausdorff. Moreover, in each of these topologies, a point (α, n) of Y is either isolated or has a neighbourhood homeomorphic to the ordinal space $\omega^m + 1$, for some $m \leq n$. Therefore, for each $i \in 3$, $(Y, \mathcal{T}_i)$ is regular, first countable, locally countable, locally compact, zero-dimensional and locally metrizable.

Claim 1. *For each $i \in 3$ and all $\alpha \in \kappa^+$, the subspace $Y_\alpha = \alpha \times \omega$ of $(Y, \mathcal{T}_i)$ is metrizable.*

Proof of Claim 1. Fix $i \in 3$. The proof is by induction so assume that Y_β is metrizable for all $\beta \in \alpha$.

Since E is a non-reflecting stationary set, each of whose elements has countable cofinality, if α is a limit ordinal (either in E or not), or $\alpha \leq \omega_1$, then there is a sequence $\{\alpha_\gamma : \gamma \in \theta \leq \alpha\}$, which is both closed, cofinal in α , and disjoint from E . But then

$$\{(\alpha_\gamma, \alpha_{\gamma+1}) \times \omega : \gamma \in \theta\} \cup \bigcup \{ \{\alpha_\gamma\} \times \omega : \gamma \in \theta \}$$

partitions Y_α into disjoint, clopen, metrizable subsets.

Now suppose that $\alpha = \beta + 1$. Without loss of generality, we may assume that β is a limit ordinal. If β is not in E , then the two sets Y_β and $\{(\beta, n) : n \in \omega\}$ partition Y_α into disjoint, clopen, metrizable sets, and we are done. Assume that β is an element of E . By construction, $\{N_j\}_{j \in \omega}$, where $N_j = N_i((\beta, j), 1) \in \mathcal{N}_i(\beta, j)$, forms a disjoint collection of clopen, metrizable subsets of Y_α . Furthermore, by $\dagger$ and $\ddagger$, if x_j is any point of N_j , then the set $\{\pi(x_j)\}_{j \in \omega}$ forms an ω -sequence, cofinal in β (though not necessarily indexed in increasing order), so the only possible limit point of the sequence $\{x_j\}_{j \in \omega}$ is (β, k) for some k in ω , which is impossible. Therefore $\{N_j\}_{j \in \omega}$ is a discrete collection of countable, clopen sets. But now $N = \bigcup_{j \in \omega} N_j$ and $Z = Y_\alpha - N$ partition Y_α into disjoint, clopen, metrizable subspaces, and again Y_α is metrizable.

Claim 2. *Fix $i \in 3$. If H is a subset of $(Y, \mathcal{T}_i)$ of size κ^+ , then H has a limit point, and, if C and D are closed subsets of $(Y, \mathcal{T}_i)$, both of size κ^+ , then C and D are not disjoint.*

Proof of Claim 2. For any subset A of Y let $A(n) = A \cap (\kappa^+ \times \{n\})$

Suppose that H has size κ^+ , then $H(n)$ also has size κ^+ for some $n \in \omega$. By $\clubsuit_{\kappa^+ \times \kappa^+}(E, 2)$, $B(\alpha, n) \times \{n\}$, is contained in $H(n)$, for some α in E , so H has $(\alpha, n + 1)$ as a limit point in $\kappa^+ \times \{n + 1\}$. In fact, since κ^+ is a regular cardinal, H has κ^+ limit points in $\kappa^+ \times \{n + 1\}$.

Now let C and D be closed subsets of $(Y, \mathcal{T}_i)$ of cardinality κ^+ . From the previous paragraph it is clear that $|C(n)| = |D(n)| = \kappa^+$, for some n . By $\clubsuit_{\kappa^+ \times \kappa^+}(E, 2)$, there is an α in E for which both $B(\alpha, 0, n) \times \{n\}$ is a subset of $C(n)$, and $B(\alpha, 1, n) \times \{n\}$ is a subset of $D(n)$, so C and D have a common limit point.

For each $i \in 3$, the (strong) collectionwise normality of $(Y, \mathcal{T}_i)$ is immediate from Claims 1 and 2: $\mathcal{D}$ be discrete collection of closed sets. By Claim 2, $\mathcal{D}$ has size less than κ^+ and there is some successor α such that Y_α contains all but at most one of the sets in $\mathcal{D}$. Since Y_α is clopen and metrizable we are done.

The strong zero-dimensionality of $(Y, \mathcal{T}_i)$ also follows from Claims 1 and 2: Suppose that A and B are subsets of Y which are completely separated by the function $f : (Y, \mathcal{T}_i) \rightarrow [0, 1]$ in that $f''A = \{0\}$ and $f''B = \{1\}$. The sets $f^{-1}''[0, 1/4]$ and $f^{-1}''[3/4, 1]$ are disjoint, closed sets containing A and B respectively, so, as above, there is a successor α such that Y_α contains A , say. Y_α is a metrizable, locally compact, zero-dimensional subspace of Y and is, therefore, strongly zero-dimensional (by 6.2.10 [E]).

Claim 3. $(Y, \mathcal{T}_i)$ is countably paracompact for each $i \in 3$.

Proof of Claim 3. Fix $i \in 3$. Since $(Y, \mathcal{T}_i)$ is normal it suffices to show that, for every decreasing sequence of closed subsets $\{D_n\}_{n \in \omega}$ of $(Y, \mathcal{T}_i)$ with empty intersection, there is a sequence of open subsets $\{U_n\}_{n \in \omega}$ with empty intersection such that U_n contains D_n .

Let $\{D_n\}_{n \in \omega}$ be such a sequence of closed sets. Suppose that each D_n has size κ^+ , then, with the notation used above, Claim 2 implies that $D_n(k)$ has size κ^+ for all k greater than some $k_n \in \omega$. By relabelling and adding repetitions if necessary, we may assume that $D_n(n)$ has size κ^+ for all n larger than some $n_0 > 0$. Now, by $\clubsuit_{\kappa^+}^\cap(E, 2)$,

$$S = \bigcap_{n \in \omega} \{\alpha \in E : T_{\alpha, i, n} \subseteq D_n(n), i \in 2\}$$

is a stationary set, and therefore non-empty. By the construction of the topology $\mathcal{T}_i$, if α is in S , then $(\alpha, 0)$ is in D_n for all $n \in \omega$, and so $\bigcap D_n$ is not empty—a contradiction.

Pick n_0 such that $|D_n| \leq \kappa$ for all $n \geq n_0$. By Claim 1 there is a successor α such that D_n is a subset of Y_α for $n \geq n_0$. The claim follows since Y_α is clopen and metrizable we are done.

Claim 4. For $i, j \in 2$, $(Y, \mathcal{T}_i) \times (Y, \mathcal{T}_j)$ is normal.

Proof of Claim 4. Let C and D be disjoint closed subsets of $(Y, \mathcal{T}_i) \times (Y, \mathcal{T}_j)$, and recall that $\Pi : (\kappa^+ \times \omega)^2 \rightarrow \kappa^+ \times \kappa^+$ is the natural projection.

Suppose that both $\Pi''C$ and $\Pi''D$ are 2-unbounded in $\kappa^+ \times \kappa^+$. There are integers $m, n, j, k \in \omega$ such that $C_{n,k} = \{(\gamma, \delta) : ((\gamma, n), (\delta, k)) \in C\}$ and $D_{m,j} = \{(\gamma, \delta) : ((\gamma, m), (\delta, j)) \in D\}$ are both 2-unbounded. Let $s = n + m + j + k + 1$, so that s is strictly greater than n, m, j and k . By $\clubsuit_{\kappa^+ \times \kappa^+}(E, 2)$, there is some α in E such that $S_{\alpha, 0}$ is a subset of $C_{n,k}$ and $S_{\alpha, 1}$ is a subset of $D_{m,j}$. By the definition of the sequence $B(\alpha, 0, s)$

$$C_{n,k} \cap (B_{\alpha, 0, s} \times B(\alpha, 0, s))$$

is infinite and cofinal in (α, α) . By the definition of the topologies $\mathcal{T}_i$ and $\mathcal{T}_j$,

$$C \cap ((B(\alpha, 0, s) \times \{n\}) \times (B_{\alpha, 0, s} \times \{k\}))$$

is cofinal in $((\alpha, s), (\alpha, s))$, which is therefore a limit point of C . Similarly $((\alpha, s), (\alpha, s))$ is a limit point of D , and C and D are not disjoint.

So suppose that $\Pi''C$ is not 2-unbounded. Choose γ not in E such that C is a subset of

$$K = ((\gamma \times \omega) \times (\kappa^+ \times \omega)) \cup ((\kappa^+ \times \omega) \times (\gamma \times \omega))$$

. Since γ is not in E , K is a clopen subset of $(Y, \mathcal{T}_i) \times (Y, \mathcal{T}_j)$. Since E is a non-reflecting stationary set, there is a club set H of γ , enumerated as $\{\gamma_\lambda : \lambda \in \theta \leq \gamma\}$, which misses E and such that $G_\lambda = \{\alpha : \gamma_\lambda < \alpha < \gamma_{\lambda+1}\}$ is countable. Now $\{\{\gamma_\lambda\} \times \omega\}_{\lambda \in \theta} \cup \{G_\lambda \times \omega\}_{\lambda \in \theta}$ partitions Y_γ into countable, metrizable, $\mathcal{T}_i$ -clopen subsets of Y , for $i = 0$ or 1 . Lemma 2.8 of [B1] states that, for normal, countably paracompact space X and a countable metric space M , $X \times M$ is normal. It is

easy to see, then, that K is normal. Since K is clopen, $(Y, \mathcal{T}_i) \times (Y, \mathcal{T}_j)$ is now, itself, seen to be normal—proving the claim.

The proof that $(Y, \mathcal{T}_i) \times (Y, \mathcal{T}_j)$ is collectionwise normal is similar

Now, let X be the disjoint topological sum of $(Y, \mathcal{T}_0)$ and $(Y, \mathcal{T}_1)$. From the above, it is clear that X satisfies the properties listed in the statement of the Theorem 1, except that it remains to show that X^2 is not countably paracompact:

Claim 5. *The closed subspace $\Delta = \{((\alpha, n), (\alpha, n)) : \alpha \in \kappa^+, n \in \omega\}$ of $(Y, \mathcal{T}_0) \times (Y, \mathcal{T}_1)$ is not countably metacompact.*

Proof of Claim 5. Let $D_n = \{((\alpha, j), (\alpha, j)) : \alpha \in \kappa^+, j \geq n\}$, and let U_n be any open subset of Δ containing D_n . $\{D_n\}_{n \in \omega}$ is a decreasing sequence of closed subsets of Δ with empty intersection, so it is enough to show that $\bigcap U_n$ is non-empty.

Notice that, since the sequences $C(\alpha, 0)$ and $C(\alpha, 1)$ are disjoint, the point $((\alpha, 0), (\alpha, 0))$ is isolated for each $\alpha \in \kappa^+$ (by hypothesis (7)). However, if α is a limit in E , then $(\alpha, n+1)$ is both a $\mathcal{T}_0$ - and a $\mathcal{T}_1$ -limit of the sequence $B(\alpha, n) \times \{n\}$. So, as $\{B(\alpha, i, n)\}_{i \in 2}^{\alpha \in E \cap \text{LIM}}$ is a $\clubsuit_{\kappa^+}(E, 2)$ -sequence, the proof of Claim 2 is, almost verbatim, a proof of:

* If H is a subset of Δ of size κ^+ , then H has a limit point in Δ , and, if C and D are closed subsets of Δ , both of size κ^+ , then C and D are not disjoint.

D_n and $\Delta - U_n$ are disjoint closed subsets. D_n has cardinality κ^+ , so, by *, $|\Delta - U_n| \leq \kappa$. Hence $|\bigcup_{n \in \omega} U_n| = \kappa^+$ and in particular Δ is not countably metacompact. This completes the proof of the Theorem. $\square$

REFERENCES

- [B1] Bešlagić A., *A Dowker product*, Trans. Amer. Math. Soc. **292** (1985), 519-530.
- [B2] Bešlagić A., *Another Dowker product*, Top. Appl. **36** (1990), 553-264.
- [B3] Bešlagić A., *Yet another Dowker product*, preprint.
- [Dv] Devlin K., *Constructability*, Springer Verlag, Berlin, 1984.
- [DJ] Dodd A.J. and Jensen R., *The covering lemma for K* , Ann. Math. Logic **22** (1982), 1-30.
- [Dk] Dowker C.H., *On countably paracompact spaces*, Canad. J. Math. **3** (1951), 219-224.
- [E] Engelking R., *General Topology*, Heldermann Verlag, Berlin, 1989.
- [F] Fleissner W.G., *The normal Moore space conjecture and large cardinals*, in: [KV], 733-760.
- [G] Good C., *Large cardinals and small Dowker spaces*, to appear in Proc. Amer. Math. Soc..
- [K] Kunen K., *Set Theory, An Introduction to Independence Proofs*, North Holland, Amsterdam, 1984.
- [KV] Kunen K. and Vaughan J.E. eds., *Handbook of Set-Theoretic Topology*, North Holland, Amsterdam, 1984.
- [R] Rudin M.E., *Dowker spaces*, in: [KV], 761-780.
- [RS] Rudin M.E and Starbird M., *Products with a metric factor*, Gen. Top. Appl. **5**, 235-248.

Wadham College, Oxford, UK.

E-mail: cgood@maths.ox.ac.uk