

Control of Fusion and Solubility in Fusion Systems

David A. Craven

March 2009

Abstract

In this article, we consider control of fusion, quotients, and p -soluble fusion systems. For control of fusion, we prove the three main theorems in the literature in a new, largely elementary way, significantly shortening their proofs. To prove one of these, and a theorem of Aschbacher that the product of strongly closed subgroups is strongly closed, we produce a consolidated treatment of quotients, collating and expanding the constructions previously available; we include analogues of the isomorphism theorems for fusion systems. We move on to p -soluble fusion systems, and prove that they are constrained, allowing us to effectively characterize fusion systems of p -soluble groups. This leads us to recast Thompson Factorization for $Qd(p)$ -free fusion systems, and consider it for more general fusion systems.

1 Introduction

The theory of fusion systems, which was started by Puig but lay unpublished for decades, attempts to formalize the concepts of local finite group theory, and extend such results to the sphere of blocks of finite groups. It has attracted considerable and growing attention over the course of the last decade, but several basic questions remain unresolved in the field. In addition, some of the results in the literature permit considerable shortenings in their proofs; the improved exposition will make clear the reasons behind the results. In Section 2 we will define the terms that we use in this introduction.

The purpose of this article is three-fold: to provide elementary proofs of some theorems in control of fusion, to provide a complete treatment of the theory of morphisms and quotients, and to examine p -soluble fusion systems. Theorems A to D have been proved in the literature, but here we provide elementary proofs. Sections 5 and 6 is a thorough treatment of morphisms and quotients, including analogues for fusion systems of the isomorphism theorems. The final two sections give the first treatment of p -soluble fusion systems available in the literature. We prove directly that they are constrained (this was proved via the generalized Fitting subsystem in [3]), and then determine those saturated fusion systems that come from p -soluble groups using a purely internal characterization.

Theorem A (Stancu [13, Proposition 6.2]) Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a subgroup of P . Then $\mathcal{F} = N_{\mathcal{F}}(Q)$ if and only if $\mathcal{F}_Q(Q)$ is a weakly normal subsystem of \mathcal{F} .

The proof given here is the first proof that does not rely on a non-trivial result: in [13], this result was given as a corollary to what is here Theorem C; in [11], this result was a corollary of what is here Theorem B; and in [3], the proof of this result requires the non-trivial fact that a constrained fusion system is the fusion system of a finite group (see [5, Proposition C]). Here we will prove it by elementary means, using the extension of the Frattini argument to fusion systems, as proved in [4].

Using Theorem A, we derive Theorem B as a consequence.

Theorem B (Linckelmann [11, Theorem 9.1]) Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let \mathcal{E} be a weakly normal subsystem, on a subgroup Q of P . Suppose that $\mathcal{E} = N_{\mathcal{E}}(R)$ for some subgroup R , and let S be the subgroup of Q generated by all \mathcal{F} -conjugates of R . Then $\mathcal{F} = N_{\mathcal{F}}(S)$.

Finally, using a result of Aschbacher's on an equivalent condition to fusion being controlled by a subgroup, we derive Stancu's main theorem of [13].

Theorem C (Stancu [13, Theorem 4.8]) Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a subgroup of P . Then $\mathcal{F} = N_{\mathcal{F}}(Q)$ if and only if Q is strongly \mathcal{F} -closed, and there is a central series

$$1 = Q_0 \leq Q_1 \leq \cdots \leq Q_n = Q$$

with each of the Q_i weakly \mathcal{F} -closed.

Moving away from control of fusion – i.e., a subgroup Q such that $\mathcal{F} = N_{\mathcal{F}}(Q)$ – we will reprove a theorem of Aschbacher from [3].

Theorem D (Aschbacher [3, Theorem 2]) Let \mathcal{F} be a fusion system. Then the product of two strongly \mathcal{F} -closed subgroups is strongly \mathcal{F} -closed.

The proof of this result in [3] is quite difficult, and requires considerable preliminaries. Here we will produce an extremely short and trivial proof of the result. However, in order to do so, we must consider factor systems, a subject that has not yet been fully understood, with differing accounts of it in the literature (see [12], [13], [11], and [4] for four, all different to various extents, approaches). Here we combine the various approaches to produce the first comprehensive treatment of the subject in Section 5, including the versions for fusion systems of the isomorphism theorems.

Theorem E Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q and R be strongly \mathcal{F} -closed subgroups with $Q \leq R$. Let \mathcal{E} be a saturated subsystem of \mathcal{F} , defined on the subgroup S

of P . Write $\mathcal{E}Q/Q$ for the image of \mathcal{E} in \mathcal{F}/Q . We have the isomorphisms

$$\mathcal{E}Q/Q \cong \mathcal{E}/(S \cap Q) \quad \text{and} \quad (\mathcal{F}/R)/(Q/R) \cong \mathcal{F}/Q.$$

We end with a discussion of p -soluble systems, which has not been considered in the literature before, although there are some unpublished notes on the subject by various authors. Broadly speaking, a fusion system is called *p-soluble* if repeated quotienting out by $O_p(\mathcal{F})$ eventually exhausts the group: a formal definition will be given in Section 7. The main result in this section is the following. (In the context of the generalized Fitting subsystem, this is also proved in [3].)

Theorem F (Aschbacher) Let \mathcal{F} be a saturated fusion system on the finite p -group P . If \mathcal{F} is p -soluble, then

$$C_P(O_p(\mathcal{F})) \leq O_p(\mathcal{F}).$$

In particular, \mathcal{F} is a constrained fusion system, so is the fusion system of a finite group.

In light of the fact that saturated subsystems of p -soluble systems are p -soluble, this gives an easy proof of the known fact that all fusion systems of p -blocks of p -soluble groups are fusion systems of finite groups.

In the final section, we consider the fusion systems of p -soluble groups; these can be effectively characterized internally.

Theorem G Let \mathcal{F} be a saturated fusion system, and write $Q = O_p(\mathcal{F})$. Then \mathcal{F} is the fusion system of a p -soluble group if and only if \mathcal{F} is constrained and $\text{Aut}_{\mathcal{F}}(Q)$ is p -soluble.

Using this, we make some remarks on extending Thompson Factorization to constrained fusion systems, and how such results differ from the versions for groups.

We begin in the next section with the definitions and concepts from fusion systems that we need. Because the notation and terminology has not yet crystallized, we will define everything to avoid ambiguity; this section contains Alperin's fusion theorem. After a preliminary section we set about proving the first two theorems given above. In the final sections, from Section 5 onwards, we introduce and study factor systems, prove Theorems C and D, and then consider p -soluble systems. Note that our maps will be composed from left to right.

2 Fusion Systems

Here we collect the very basic concepts in the field of fusion systems. We begin by defining fusion systems: for more background see [11] (note that fusion systems there are saturated fusion systems here) or [7]. If g is an element of a group G , denote by c_g the map induced by conjugation by g .

Definition 2.1 Let P be a finite p -group. A *fusion system* \mathcal{F} on P is a category, whose objects are the set of all subgroups of P , and whose morphisms $\text{Hom}_{\mathcal{F}}(Q, R)$ are injective homomorphisms

$Q \rightarrow R$ (with composition in the category the usual composition), satisfying the following three axioms:

- (i) for each $g \in P$ with $Q^g \leq R$, the associated conjugation map $c_g : Q \rightarrow R$ is in $\text{Hom}_{\mathcal{F}}(Q, R)$;
- (ii) for each $\phi \in \text{Hom}_{\mathcal{F}}(Q, R)$, the isomorphism $Q \rightarrow Q\phi$ lies in $\text{Hom}_{\mathcal{F}}(Q, Q\phi)$; and
- (iii) if $\phi \in \text{Hom}_{\mathcal{F}}(Q, R)$ is an isomorphism, then its inverse $\phi^{-1} : R \rightarrow Q$ lies in $\text{Hom}_{\mathcal{F}}(R, Q)$.

We write $\text{Aut}_{\mathcal{F}}(Q)$ for the set (in fact group) $\text{Hom}_{\mathcal{F}}(Q, Q)$.

Observe that it is a consequence of the first axiom that if Q is a subgroup of R then the inclusion map from Q into R is a morphism in \mathcal{F} . If G is a finite group with Sylow p -subgroup P , we write $\mathcal{F}_P(G)$ for the fusion system obtained from conjugation by elements of G on the subgroups of P . We also note the *universal fusion system* $\mathcal{U}(P)$ on a finite p -group P , where $\text{Hom}_{\mathcal{U}(P)}(Q, R)$ is the set of *all* injective homomorphisms from Q to R . Obviously any fusion system is a subsystem of the universal fusion system.

Definition 2.2 Let P be a finite p -group, and let Q be a subgroup of P . Let \mathcal{F} be a fusion system on P . We say that Q is *fully \mathcal{F} -normalized* if, whenever $\phi : Q \rightarrow R$ is an isomorphism in \mathcal{F} , we have that $|\text{N}_P(Q)| \geq |\text{N}_P(R)|$. If the fusion system involved is clear from the context, we will simply say fully normalized.

Definition 2.3 Let P be a finite p -group, and let \mathcal{F} be a fusion system on P . We say that \mathcal{F} is *saturated* if

- (i) $\text{Aut}_P(P)$ is a Sylow p -subgroup of $\text{Aut}_{\mathcal{F}}(P)$, and
- (ii) every morphism $\phi : Q \rightarrow P$ in \mathcal{F} such that $Q\phi$ is fully normalized extends to a morphism $\bar{\phi} : N_{\phi} \rightarrow P$, where

$$N_{\phi} = \{x \in \text{N}_P(Q) : \text{there exists } y \in \text{N}_P(Q\phi) \text{ such that } (g^x)\phi = (g\phi)^y \text{ for all } g \in Q\}.$$

Note that, if $\phi : Q \rightarrow R$ is an \mathcal{F} -isomorphism, then N_{ϕ} is the inverse image under the map $\delta_Q : \text{N}_P(Q) \rightarrow \text{Aut}(Q)$ of the subgroup $\text{Aut}_P(Q) \cap \phi \text{Aut}_P(R)\phi^{-1}$, and that $QC_P(Q)$ is always contained within N_{ϕ} .

Definition 2.4 Let \mathcal{F} be a fusion system on the finite p -group P . Let Q be a subgroup of P , and let K be a subgroup of $\text{Aut}(Q)$.

- (i) The subgroup $\text{N}_P^K(Q)$, the *K -normalizer* of Q in P , is the set of all $g \in \text{N}_P(Q)$ such that g induces by conjugation an automorphism of Q that lies in K . Write $\text{Aut}_P(Q)$ for the subgroup of $\text{Aut}(Q)$ of automorphisms induced by elements of P , and $\text{Aut}_P^K(Q) = K \cap \text{Aut}_P(Q)$.

- (ii) The fusion system $N_{\mathcal{F}}^K(Q)$ is the category whose objects are all subgroups of $N_P^K(Q)$, and whose morphisms $\text{Hom}_{N_{\mathcal{F}}^K(Q)}(R, S)$ are

$$\{\phi \in \text{Hom}_{\mathcal{F}}(R, S) : \text{there is } \bar{\phi} \in \text{Hom}_{\mathcal{F}}(QR, QS) \text{ with } \bar{\phi}|_R = \phi \text{ and } \bar{\phi}|_Q \in K\}.$$

The fusion system $N_{\mathcal{F}}^K(Q)$ is called the *K-normalizer subsystem* of Q in \mathcal{F} .

Of particular interest are the cases $K = \text{Aut}(Q)$, in which case we write $N_{\mathcal{F}}(Q)$ and call this the *normalizer* of Q in \mathcal{F} , and $K = 1$, in which case we write $C_{\mathcal{F}}(Q)$ and call this the *centralizer* of Q in \mathcal{F} .

If Q is fully normalized and $K \trianglelefteq \text{Aut}_{\mathcal{F}}(Q)$ then $N_{\mathcal{F}}^K(Q)$ is saturated (see [11, Theorem 3.6]). If Q is a subgroup of P , we are interested in the case where $\mathcal{F} = N_{\mathcal{F}}(Q)$, in which case we say that the subgroup is *normal* in \mathcal{F} . Theorems A and C give equivalent conditions for a subgroup to be normal in \mathcal{F} . We need a few more of these results in the course of this article, but in order to state the first one in Section 3 we need the concept of \mathcal{F} -centric and \mathcal{F} -radical subgroups.

Let \mathcal{F} be a fusion system on a finite p -group P , and let Q be a subgroup of P . We say that Q is *\mathcal{F} -centric* if, whenever R is \mathcal{F} -isomorphic to Q , then R contains its centralizer (or equivalently, $C_P(R) = Z(R)$). We say that Q is *\mathcal{F} -radical* if $O_p(\text{Aut}_{\mathcal{F}}(Q)) = \text{Inn}(Q)$. A fusion system \mathcal{F} is *constrained* if it contains a normal, \mathcal{F} -centric subgroup. The fundamental theorem on such systems is the following.

Theorem 2.5 ([5, Proposition C]) Let \mathcal{F} be a constrained saturated fusion system on a finite p -group P . Then there is a unique finite group G such that

- (i) $O_{p'}(G) = 1$,
- (ii) $C_G(O_p(G)) \leq O_p(G)$, and
- (iii) $\mathcal{F} = \mathcal{F}_P(G)$.

Using the concept of centric and radical subgroups, we also have a version for fusion systems of Alperin's fusion theorem.

Theorem 2.6 (Alperin's fusion theorem) Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let $\phi : Q \rightarrow R$ be an isomorphism. Then there exist

- (i) a sequence of \mathcal{F} -isomorphic subgroups $Q = Q_0, Q_1, \dots, Q_n = R$,
- (ii) a sequence S_1, S_2, \dots, S_n of fully normalized, \mathcal{F} -radical, \mathcal{F} -centric subgroups, with $Q_{i-1}, Q_i \leq S_i$, and
- (iii) a sequence of \mathcal{F} -automorphisms ϕ_i of S_i with $Q_{i-1}\phi_i = Q_i$,

such that

$$(\phi_1\phi_2 \dots \phi_n)|_Q = \phi.$$

A proof may be found in, for example, [7, Theorem A.10], and a stronger version is in [11].

Now we define two types of subgroups: a subgroup Q is called *weakly \mathcal{F} -closed* if Q is only \mathcal{F} -isomorphic to itself, and it is called *strongly \mathcal{F} -closed* if whenever any \mathcal{F} -morphism has domain a subgroup of Q , it has image a subgroup of Q .

Definition 2.7 Let \mathcal{F} be a fusion system on a finite p -group P , and let Q be a subgroup of P . A *subsystem* \mathcal{E} of \mathcal{F} , on Q , is a subcategory of \mathcal{F} that is a fusion system on Q . We say that \mathcal{E} is *\mathcal{F} -invariant* if Q is strongly \mathcal{F} -closed and, for each $R \leq S \leq Q$, $\phi \in \text{Hom}_{\mathcal{E}}(R, S)$, and $\psi \in \text{Hom}_{\mathcal{F}}(S, P)$, we have that $\psi^{-1}\phi\psi$ is a morphism in $\text{Hom}_{\mathcal{E}}(R\psi, Q)$. If, in addition, \mathcal{E} is saturated, we say that \mathcal{E} is *weakly normal* in \mathcal{F} . We denote weak normality by $\mathcal{E} \prec \mathcal{F}$. A saturated fusion system is *simple* if it contains no non-trivial, proper, weakly normal subsystems.

An *automorphism* of \mathcal{F} is an automorphism α of P such that, if $\phi : Q \rightarrow R$ is a morphism in \mathcal{F} , then the induced map $\phi' : Q\alpha \rightarrow R\alpha$ also lies in \mathcal{F} . The set of all automorphisms of \mathcal{F} is denoted by $\text{Aut}(\mathcal{F})$. If \mathcal{F} is a saturated fusion system and \mathcal{E} is a weakly normal subsystem, then \mathcal{E} is said to be *characteristic* if $\mathcal{E}\phi = \mathcal{E}$ for all $\phi \in \text{Aut}(\mathcal{F})$. We denote this by $\mathcal{E} \text{ char } \mathcal{F}$.

We have called this definition ‘weakly normal’, because (in what is becoming standard terminology, at least at this point) ‘normal’ refers to the definition of normality given by Aschbacher in [4]. In [4], for a weakly normal subsystem \mathcal{E} , defined on Q , to be *normal* in \mathcal{F} , defined on P , it is required that every automorphism $\phi \in \text{Aut}_{\mathcal{E}}(Q)$ extends to an automorphism $\bar{\phi} \in \text{Aut}_{\mathcal{F}}(Q C_P(Q))$ with $\bar{\phi}$ acting trivially on $Q C_P(Q)/Q$. This extra piece of information enables Aschbacher to develop [3] [4] a local theory of fusion systems, as well as prove the existence of normal subsystems acting like the intersection of two normal subsystems, and the central product of two normal subsystems. The former of these is not known to exist for weakly normal subsystems, and the latter is known not to exist. Here we will focus on weakly normal subsystems, because they share many properties of normal subsystems, and also they are the easiest to work with in situations of control of fusion and p -solubility. (For example, it is easy to show that if \mathcal{F} is a saturated fusion system on P , and Q is a subgroup of P , then $\mathcal{F}_Q(Q)$ is a weakly normal subsystem of \mathcal{F} if and only if it is a normal subsystem of \mathcal{F} .)

Because we need to use a variant of the Frattini argument for fusion systems, we need the definition of an \mathcal{F} -Frattini subsystem.

Definition 2.8 ([4, Section 3]) Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let \mathcal{E} be a subsystem of \mathcal{F} , on the subgroup Q . We say that \mathcal{E} is *\mathcal{F} -Frattini* if, whenever $R \leq Q$ and $\phi : R \rightarrow P$ is a morphism in \mathcal{F} , there exist morphisms $\alpha \in \text{Aut}_{\mathcal{F}}(Q)$ and $\beta \in \text{Hom}_{\mathcal{E}}(R\alpha, P)$ such that $\phi = \alpha\beta$.

3 Preliminaries

Our first result tells us that we may apply the Frattini argument whenever \mathcal{E} is a weakly normal subsystem of \mathcal{F} , and in fact gives us an equivalent condition to weak normality of subsystems.

Theorem 3.1 (Puig [12, Proposition 6.6]) Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let \mathcal{E} be a subsystem on a strongly \mathcal{F} -closed subgroup Q . Then \mathcal{E} is \mathcal{F} -invariant if and only if $\text{Aut}_{\mathcal{F}}(Q) \leq \text{Aut}(\mathcal{E})$ and \mathcal{E} is \mathcal{F} -Frattini.

We also need the equivalent for fusion systems of the statement for groups that if $K \text{ char } H \trianglelefteq G$ then $K \trianglelefteq G$.

Proposition 3.2 (Aschbacher [3, 7.4]) Let \mathcal{F} be a saturated fusion system on a finite p -group P and suppose that $\mathcal{E} \prec \mathcal{E}' \prec \mathcal{F}$, where \mathcal{E}' is defined on the subgroup Q and \mathcal{E} is defined on R . Suppose that all \mathcal{F} -automorphisms of Q induce automorphisms on \mathcal{E} , that is if $\alpha \in \text{Aut}_{\mathcal{F}}(Q)$ then $\alpha|_R \in \text{Aut}(\mathcal{E})$. Then $\mathcal{E} \prec \mathcal{F}$. In particular, if $\mathcal{E} \text{ char } \mathcal{E}' \prec \mathcal{F}$, then $\mathcal{E} \prec \mathcal{F}$.

Proof: Suppose that $\mathcal{E} \text{ char } \mathcal{E}'$. Then $\text{Aut}_{\mathcal{F}}(Q) \leq \text{Aut}(\mathcal{E}')$ by Theorem 3.1, and so since \mathcal{E} is $\text{Aut}(\mathcal{E}')$ -invariant, the second statement follows from the first. Let R be the subgroup of Q on which \mathcal{E} acts.

If $S \leq R$ and $\phi : S \rightarrow P$ is any map in \mathcal{F} , then since Q is strongly \mathcal{F} -closed, we have that $\text{im } \phi \leq Q$. Since $\mathcal{E}' \prec \mathcal{F}$, it is \mathcal{F} -Frattini by Theorem 3.1, and so $\phi = \alpha\beta$, where $\alpha \in \text{Aut}_{\mathcal{F}}(Q)$ and β is a morphism in \mathcal{E}' . Since all \mathcal{F} -automorphisms of Q induce automorphisms on \mathcal{E} , we see that $S\alpha \leq R$, and as R is strongly \mathcal{E}' -closed, $S\phi = (S\alpha)\beta \leq R$. In particular, R is strongly \mathcal{F} -closed as $S\phi \leq R$.

We will use Theorem 3.1 again: since $\mathcal{E} \prec \mathcal{E}'$, \mathcal{E} is \mathcal{E}' -Frattini, so $\beta = \gamma\delta$, where $\gamma \in \text{Aut}_{\mathcal{E}'}(R)$ and δ is a morphism in \mathcal{E}' . Then $\phi = \alpha\gamma\delta$. We claim that this decomposition proves that \mathcal{E} is \mathcal{F} -Frattini; since $\alpha \in \text{Aut}_{\mathcal{F}}(Q)$ and R is strongly \mathcal{F} -closed, $\alpha|_R$ is an automorphism, and hence $\alpha\gamma \in \text{Aut}_{\mathcal{F}}(R)$. Since $\delta \in \text{Hom}_{\mathcal{E}}(S\alpha\gamma, R)$, this gives the correct decomposition of ϕ .

If $S = R$, then we perform the same decomposition to get that $\phi \in \text{Aut}_{\mathcal{F}}(R)$ may be written as $\phi = \alpha\gamma$, where $\alpha \in \text{Aut}_{\mathcal{F}}(Q)$ and $\gamma \in \text{Aut}_{\mathcal{E}'}(R)$. Because all elements of $\text{Aut}_{\mathcal{F}}(Q)$ induce automorphisms on \mathcal{E} by hypothesis, and $\text{Aut}_{\mathcal{E}'}(R) \leq \text{Aut}(\mathcal{E})$ by Theorem 3.1, we see that $\text{Aut}_{\mathcal{F}}(R) \leq \text{Aut}(\mathcal{E})$, and so $\mathcal{E} \prec \mathcal{F}$ by Theorem 3.1. \square

We also need two equivalent conditions for a subgroup to be normal in a fusion system.

Proposition 3.3 ([5, Proposition 1.6]) Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a strongly \mathcal{F} -closed subgroup of P . Then $\mathcal{F} = \text{N}_{\mathcal{F}}(Q)$ if and only if Q is contained in every fully normalized, \mathcal{F} -centric, \mathcal{F} -radical subgroup.

Proposition 3.4 (Aschbacher [2, (3.7)]) Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a subgroup of P . Then $\mathcal{F} = \mathcal{N}_{\mathcal{F}}(Q)$ if and only if there exists a central series

$$1 = Q_0 \leq Q_1 \leq \cdots \leq Q_n = Q$$

for Q , all of whose terms are strongly \mathcal{F} -closed.

Proof: Suppose that $\mathcal{F} = \mathcal{N}_{\mathcal{F}}(Q)$, and let $Q_i = Z_i(Q)$. (The subgroup $Z_i(Q)$ is the preimage in Q of $Z(Q/Z_{i-1}(Q))$, where $Z_0(Q) = 1$.) We need to show that every Q_i is strongly \mathcal{F} -closed, and we are done. Let $R \leq Q_i$, and $\phi \in \text{Hom}_{\mathcal{F}}(R, P)$. Since Q is normal in \mathcal{F} and $R \leq Q$, it follows that ϕ extends to $\bar{\phi} \in \text{Hom}_{\mathcal{F}}(Q, P)$ that acts like an automorphism on Q (using that Q is strongly \mathcal{F} -closed). Since by construction $Q_i \text{ char } Q$, we see that $\text{im } \phi \leq Q_i$, as needed.

Now suppose that (ii) holds, and let T be any fully \mathcal{F} -normalized, \mathcal{F} -radical, \mathcal{F} -centric subgroup. If we can show that in this case $Q \leq T$, then we are done, since then Q is contained in every \mathcal{F} -radical, \mathcal{F} -centric subgroup, and so $\mathcal{F} = \mathcal{N}_{\mathcal{F}}(Q)$ by Proposition 3.3. Choose i maximal such that $Q_i \leq T$, so that $Q_{i+1} \not\leq T$. (For a contradiction, assume that Q is not a subgroup of T .) Set $R = Q_{i+1} \cap T$ and $S = \mathcal{N}_{Q_{i+1}}(T)$; as $Q \not\leq T$, we have that $R < S$. Since S normalizes T and $Q_{i+1} \leq P$, we have that $[T, S] \leq R$, and as Q_{i+1}/Q_i is central in Q/Q_i , we see that S centralizes R/Q_i , and since each Q_{j+1}/Q_j is central in Q/Q_j , we have that S centralizes Q_{j+1}/Q_j for all $j < i$.

Each Q_i is strongly \mathcal{F} -closed, and therefore $\text{Aut}_{\mathcal{F}}(T)$ acts on $R = Q_{i+1} \cap T$ and Q_j for all $j \leq i$. Hence there is an $\text{Aut}_{\mathcal{F}}(T)$ -invariant series

$$1 = Q_0 \leq \cdots \leq Q_i \leq R \leq T,$$

with S centralizing each factor. The set of all such automorphisms of T is clearly a normal subgroup, and is a p -subgroup by [8, Corollary 5.3.3]. Therefore, $\text{Aut}_S(T)$ is contained in a normal p -subgroup of $\text{Aut}_{\mathcal{F}}(T)$, and so in particular $\text{Aut}_S(T) \leq \text{Inn}(T)$ since T is \mathcal{F} -radical. Thus $S = \mathcal{N}_{Q_{i+1}}(T) \leq TC_P(T) = T$ since T is \mathcal{F} -centric. Therefore, $Q_{i+1} \cap T \geq S > R = Q_{i+1} \cap T$, a contradiction. Hence Q is contained in every fully \mathcal{F} -normalized, \mathcal{F} -centric, \mathcal{F} -radical subgroup, and so $\mathcal{F} = \mathcal{N}_{\mathcal{F}}(Q)$, as claimed. \square

We are now in a position to prove three preliminary lemmas – the last one perhaps of independent interest – needed to prove the first three main theorems.

Lemma 3.5 (cf. [13, Proposition 6.2]) Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a subgroup such that $\mathcal{F} = \mathcal{N}_{\mathcal{F}}(Q)$. Then $\mathcal{F}_Q(Q)$ is weakly normal in \mathcal{F} .

Proof: Suppose that $\mathcal{F} = \mathcal{N}_{\mathcal{F}}(Q)$; then any morphism $\phi : R \rightarrow S$ lifts to a morphism $\bar{\phi} : QR \rightarrow QS$ that acts as an automorphism on Q and so Q is strongly \mathcal{F} -closed. Write $\mathcal{E} = \mathcal{F}_Q(Q)$; if $R \leq S \leq Q$ and $\psi : S \rightarrow P$ is a map in \mathcal{F} , then we should show that, for all $\phi = c_g$ with $g \in Q$, the map $\psi^{-1}\phi\psi$ is also in \mathcal{E} . If $\psi : S \rightarrow P$ is a map in \mathcal{F} , then since Q is strongly \mathcal{F} -closed, this is actually $\psi : S \rightarrow Q$, and since $\mathcal{F} = \mathcal{N}_{\mathcal{F}}(Q)$, the map ψ extends to an automorphism $\bar{\psi} \in \text{Aut}_{\mathcal{F}}(Q)$. Thus we

may assume that $S = Q$; similarly, we may assume that $R = Q$ as well, so suppose that $R = S = Q$. We need to show that if $g \in Q$ and $\psi \in \text{Aut}_{\mathcal{F}}(Q)$, then $\psi^{-1}c_g\psi \in \mathcal{F}_Q(Q)$, but it is well known that $\psi^{-1}c_g\psi = c_{g\psi} \in \mathcal{F}_Q(Q)$, and we get the result. \square

The proof of the next lemma is due to Sejong Park, and replaces a less elegant proof in a preliminary version of the manuscript.

Lemma 3.6 Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a subgroup such that $\mathcal{F}_Q(Q) \prec \mathcal{F}$. Then every characteristic subgroup of Q is strongly \mathcal{F} -closed. In particular, Q has a central series each of whose terms is strongly \mathcal{F} -closed.

Proof: Let R be a characteristic subgroup of Q . Then $\mathcal{F}_R(R) \text{ char } \mathcal{F}_Q(Q) \prec \mathcal{F}$, and so $\mathcal{F}_R(R) \prec \mathcal{F}$ by Proposition 3.2; in particular, R is strongly \mathcal{F} -closed. \square

Lemma 3.7 Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a strongly \mathcal{F} -closed subgroup of P . Let Z be a central subgroup of Q that is weakly \mathcal{F} -closed. Then Z is strongly \mathcal{F} -closed.

Proof: Let $\phi : X \rightarrow Y$ be an \mathcal{F} -isomorphism with either X or Y contained in Z , and Y fully normalized. Since Q is strongly \mathcal{F} -closed, both X and Y lie inside Q . The subgroup Z is central in Q , and so $Z \leq C_P(X)$; as Y is fully normalized and \mathcal{F} is saturated, ϕ extends to a morphism $\psi : ZX \rightarrow Q$ which, as Z is weakly \mathcal{F} -closed, restricts to an automorphism $\psi|_Z : Z \rightarrow Z$. Therefore if either X or Y lies in Z , then the other also does, as required. \square

4 Theorems A and B

We begin by proving Theorem A.

Theorem A Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a subgroup of P . Then $\mathcal{F} = \text{N}_{\mathcal{F}}(Q)$ if and only if $\mathcal{F}_Q(Q) \prec \mathcal{F}$.

Proof: This follows immediately from Proposition 3.4 and Lemmas 3.5 and 3.6. \square

We now prove Theorem B, using Theorem A.

Theorem B Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let \mathcal{E} be a weakly normal subsystem, on a subgroup Q of P . Suppose that $\mathcal{E} = \text{N}_{\mathcal{E}}(R)$ for some subgroup R , and let S be the subgroup of Q generated by all \mathcal{F} -conjugates of R . Then $\mathcal{F} = \text{N}_{\mathcal{F}}(S)$.

Proof: Suppose that $\mathcal{E} = \text{N}_{\mathcal{E}}(R)$, and write S for the product of all \mathcal{F} -conjugates of R (all of which are subgroups of Q , since Q is strongly \mathcal{F} -closed). Then $\mathcal{E} = \text{N}_{\mathcal{E}}(S)$ since the product of subgroups normal in \mathcal{E} is also normal in \mathcal{E} , and if $\phi \in \text{Aut}_{\mathcal{F}}(Q)$, then ϕ leaves S invariant (as $\phi \in \text{Aut}(\mathcal{E})$ by Theorem 3.1, and clearly any automorphism of \mathcal{E} will permute the normal subgroups of \mathcal{E}), so

induces an automorphism on $\mathcal{D} = \mathcal{F}_S(S)$. (By Theorem A, $\mathcal{F}_S(S) \prec \mathcal{E}$ if and only if $\mathcal{E} = \mathcal{N}_{\mathcal{E}}(S)$.) Therefore, since $\mathcal{D} \prec \mathcal{E} \prec \mathcal{F}$, by Proposition 3.2, $\mathcal{D} \prec \mathcal{F}$. Again, by Theorem A, we get that $\mathcal{F} = \mathcal{N}_{\mathcal{F}}(S)$, as claimed. \square

Related to this, we have a proposition on the subgroup $O_p(\mathcal{F})$ of a fusion system \mathcal{F} . As we said earlier, if Q and R are normal subgroups of \mathcal{F} , then QR is as well, and hence there is a largest subgroup of P normal in \mathcal{F} ; this is denoted by $O_p(\mathcal{F})$. Denote by $\mathcal{O}_p(\mathcal{F})$ the subsystem $\mathcal{F}_Q(Q)$ of \mathcal{F} , where $Q = O_p(\mathcal{F})$. By Theorem A, $\mathcal{O}_p(\mathcal{F})$ is a weakly normal subsystem, and it is invariant under $\text{Aut}(\mathcal{F})$, so characteristic.

Proposition 4.1 Let \mathcal{F} be a saturated fusion system on a finite p -group P . If \mathcal{E} is a weakly normal subsystem of \mathcal{F} on a subgroup Q , then $O_p(\mathcal{F}) \cap Q = O_p(\mathcal{E})$. In particular, if $O_p(\mathcal{E}) \neq 1$ for some $\mathcal{E} \prec \mathcal{F}$, then $O_p(\mathcal{F}) \neq 1$.

Proof: Let $R = O_p(\mathcal{F})$; by Proposition 3.4, R possesses a central series

$$1 = R_0 \leq R_1 \leq \cdots \leq R_d = R,$$

such that each R_i is strongly \mathcal{F} -closed. We claim that $Q_i = Q \cap R_i$ is strongly \mathcal{E} -closed; in this case,

$$1 = Q_0 \leq Q_1 \leq \cdots \leq Q_d = Q \cap R$$

is a central series for $Q \cap R$, each of whose terms is strongly \mathcal{E} -closed, yielding that $Q \cap R \leq O_p(\mathcal{E})$, as required. It remains to show that Q_i is strongly \mathcal{E} -closed; however, any morphism in \mathcal{E} that originates inside $Q_i = Q \cap R_i$ must have image inside Q since \mathcal{E} is a system on Q , and must also have image in R_i since it is strongly \mathcal{F} -closed, and so Q_i is strongly \mathcal{E} -closed. Thus $Q \cap R \leq O_p(\mathcal{E})$.

On the other hand, Theorem B tells us that $O_p(\mathcal{E}) \leq O_p(\mathcal{F})$, and so we get equality. \square

In order to prove Theorems C and D, we need to understand factor systems, and we delay their proofs until after the next section.

5 Quotients

In this section we will consider morphisms of fusion systems and quotients. The treatment of these varies wildly in the literature with several opposing viewpoints and one or two errors, and it is our intention here to produce a clear description of the subject. We begin by defining a factor system.

Definition 5.1 Let \mathcal{F} be a fusion system on the finite p -group P , and let Q be a normal subgroup of P . By the *factor system* \mathcal{F}/Q , we mean the category, whose objects are all subgroups of P/Q , and such that for any two subgroups R and S containing Q , we have that $\text{Hom}_{\mathcal{F}/Q}(R/Q, S/Q)$ is the set of homomorphisms ϕ induced from the set $\text{Hom}_{\mathcal{F}}(R, S)$ such that $Q\phi = Q$.

Traditionally, in the definition above the subgroup Q is strongly \mathcal{F} -closed, but this is not necessary for the following two results. The first is easy, and a proof is omitted.

Proposition 5.2 Let \mathcal{F} be a fusion system on the finite p -group P . If Q is a normal subgroup of P , then the category \mathcal{F}/Q is a fusion system on P/Q .

Our proof of the next proposition follows [11, Theorem 6.2], and although our hypotheses are weaker, the method of proof is the same.

Proposition 5.3 Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a weakly \mathcal{F} -closed subgroup of P . Then the fusion system \mathcal{F}/Q is saturated.

Proof: All automorphisms in $\text{Aut}_{\mathcal{F}/Q}(P/Q)$ are induced from automorphisms in $\text{Aut}_{\mathcal{F}}(P)$, and so the obvious homomorphism $\text{Aut}_{\mathcal{F}}(P) \rightarrow \text{Aut}_{\mathcal{F}/Q}(P/Q)$ is surjective. The image of $\text{Aut}_P(P)$ in $\text{Aut}_{\mathcal{F}/Q}(P/Q)$ is clearly $\text{Aut}_{P/Q}(P/Q)$, so that it satisfies the first axiom of a saturated fusion system.

Suppose that $\phi \in \text{Hom}_{\mathcal{F}/Q}(R/Q, S/Q)$ is an isomorphism such that S/Q is fully \mathcal{F}/Q -normalized. We claim that S is also fully \mathcal{F} -normalized. Since $Q \leq R$, and Q is weakly \mathcal{F} -closed, for all T that are \mathcal{F} -isomorphic to R , we have that $Q \leq T$ and $Q \leq N_P(T)$. Also, $N_P(R)/Q = N_{P/Q}(R/Q)$. Therefore

$$|N_P(T)| = |N_{P/Q}(T/Q)| \cdot |Q| \leq |N_{P/Q}(S/Q)| \cdot |Q| = |N_P(S)|;$$

hence S is fully \mathcal{F} -normalized.

Now let ϕ be an automorphism of a fully \mathcal{F}/Q -normalized subgroup R/Q , and let ψ be an \mathcal{F} -automorphism of R with image ϕ in \mathcal{F}/Q . At this point we would like to prove that $N_{\psi}/Q = N_{\phi}$, but it is not necessarily true. However, there is *some* ψ with image ϕ for which $N_{\psi}/Q = N_{\phi}$, as we shall demonstrate now. Notice that $N_{\psi}/Q \leq N_{\phi}$ trivially.

Let K be the kernel of the natural map $\text{Aut}_{\mathcal{F}}(R) \rightarrow \text{Aut}(R/Q)$, a normal subgroup of $\text{Aut}_{\mathcal{F}}(R)$; then K consists of all elements of $\text{Aut}_{\mathcal{F}}(R)$ that act trivially on R/Q , and hence are sent to the identity automorphism of R/Q under the map $\mathcal{F} \rightarrow \mathcal{F}/Q$. The idea is that if $\chi \in K$, then $\chi\psi$ and ψ both have the image ϕ in \mathcal{F}/Q , so one may ‘ignore’ elements in K . We will prove that there are morphisms $\chi \in K$ and $\theta : R \rightarrow R$ such that $\psi = \chi\theta$ and θ has the property that $N_{\theta}/Q = N_{\phi}$. Since θ and ψ define the same image ϕ in \mathcal{F}/Q , we prove that ϕ extends to N_{ϕ} .

Since K is a normal subgroup of $\text{Aut}_{\mathcal{F}}(R)$ and $\text{Aut}_P(R)$ is a Sylow p -subgroup of $\text{Aut}_{\mathcal{F}}(R)$ (as R is fully normalized), we have that $\text{Aut}_P^K(R) = K \cap \text{Aut}_P(R)$ is a Sylow p -subgroup of K , and by the Frattini argument

$$\text{Aut}_{\mathcal{F}}(R) = K N_{\text{Aut}_{\mathcal{F}}(R)}(\text{Aut}_P^K(R)).$$

Since $\text{Aut}_P(R)$ normalizes $\text{Aut}_P^K(R) = \text{Aut}_P(R) \cap K$, we see that $\text{Aut}_P(R)$ is a Sylow p -subgroup of $N_{\text{Aut}_{\mathcal{F}}(R)}(\text{Aut}_P^K(R))$. Also, writing $X = N_{\text{Aut}_{\mathcal{F}}(R)}(\text{Aut}_P^K(R))$, we have

$$X/X \cap K \cong KX/K = \text{Aut}_{\mathcal{F}}(R)/K \cong \text{Aut}_{\mathcal{F}/Q}(R/Q),$$

by the second isomorphism theorem and the definition of K . Since $X = N_{\text{Aut}_{\mathcal{F}}(R)}(\text{Aut}_P^K(R))$, we may form the quotient group $X/\text{Aut}_P^K(R)$, and as $S = \text{Aut}_P^K(R)$ is a Sylow p -subgroup of K , it must be a (normal) Sylow p -subgroup of $X \cap K$; hence $(X \cap K)/S$ is a p' -group. Quotienting out by this Sylow p -subgroup, we see that

$$X/X \cap K \cong (X/S)/((X \cap K)/S) \cong \text{Aut}_{\mathcal{F}/Q}(R/Q).$$

The subgroup N_ϕ is the preimage in P/Q of the intersection $\text{Aut}_{P/Q}(R/Q) \cap \text{Aut}_{P/Q}(R/Q)^{\phi^{-1}}$; notice that both of these subgroups in the intersection are Sylow p -subgroups of $\text{Aut}_{\mathcal{F}/Q}(R/Q)$. Since $(X \cap K)/S$ is a p' -group, we see that there are two Sylow p -subgroups A/S and B/S of X/S that project onto $\text{Aut}_{P/Q}(R/Q)$ and $\text{Aut}_{P/Q}(R/Q)^{\phi^{-1}}$ respectively, and an element $Sg \in X/S$, such that $A/S \cap (B/S)^{Sg^{-1}}$ projects onto $\text{Aut}_{P/Q}(R/Q) \cap \text{Aut}_{P/Q}(R/Q)^{\phi^{-1}}$ and Sg projects onto ϕ . (For a proof, see [11, Corollary 2.13]; it essentially comes from the fact that if H is a normal p' -subgroup, then the fusion systems of G and G/H are isomorphic.)

Since S is a normal p -subgroup in $X \cap K$, we may easily lift from X/S to X , so there is some element $\theta \in X = N_{\text{Aut}_{\mathcal{F}}(R)}(\text{Aut}_P^K(R))$ such that $\text{Aut}_P(R) \cap \text{Aut}_P(R)^{\theta^{-1}}$ maps onto $\text{Aut}_{P/Q}(R/Q) \cap (\text{Aut}_{P/Q}(R/Q))^{\phi^{-1}}$. Therefore the map $N_\theta \rightarrow N_\phi$ is surjective, and so $N_\theta/Q = N_\phi$. As the map θ extends to N_θ , the map ϕ extends to N_ϕ .

It remains to deal with the general case of any map $\phi : S/Q \rightarrow R/Q$ in \mathcal{F}/Q , where R/Q is fully \mathcal{F}/Q -normalized. This lifts to a map $\psi : S \rightarrow R$ in \mathcal{F} with R fully \mathcal{F} -normalized (since it was shown at the beginning of the proof that if R/Q is fully \mathcal{F}/Q -normalized then R is fully \mathcal{F} -normalized). By [11, Lemma 2.6], there is some map $\theta : S \rightarrow R$ with $N_\theta = N_P(S)$, and so ϕ extends to N_ϕ if and only if both the image χ of θ in \mathcal{F}/Q extends to $N_\chi = N_{P/Q}(S/Q)$ and the automorphism $\chi^{-1}\phi$ of R/Q extends to $N_{\chi^{-1}\phi} = N_\phi$. The first of these claims is obvious, and the second has just been proved above, concluding the proof. \square

In the other direction, since fusion systems are categories, one may consider morphisms of fusion systems.

Definition 5.4 Let \mathcal{F} and \mathcal{E} be fusion systems on the finite p -groups P and Q respectively. A *morphism* $\Phi : \mathcal{F} \rightarrow \mathcal{E}$ of fusion systems is a pair $(\phi, \{\phi_{R,S} : R, S \leq P\})$, where $\phi : P \rightarrow Q$ is a group homomorphism, and for each $R, S \leq P$, the map $\phi_{R,S}$ is a function

$$\phi_{R,S} : \text{Hom}_{\mathcal{F}}(R, S) \rightarrow \text{Hom}_{\mathcal{E}}(R\phi, S\phi)$$

such that the corresponding map $\mathcal{F} \rightarrow \mathcal{E}$ between the categories is a functor; i.e., for any two composable \mathcal{F} -morphisms α and β , we have

$$(\alpha\beta)\Phi = \alpha\Phi\beta\Phi.$$

It is easy to see, using the functoriality of Φ , that the action of Φ on morphisms in \mathcal{F} is completely determined by the underlying group homomorphism.

Kernels of morphisms are clearly normal subgroups, since they are kernels of group homomorphisms. We have even more.

Proposition 5.5 If $\Phi : \mathcal{F} \rightarrow \mathcal{E}$ is a morphism of fusion systems, then $\ker \Phi$ is strongly \mathcal{F} -closed.

Proof: Let Q be the kernel of Φ , and let R be a subgroup of Q . We need to show that if S is \mathcal{F} -isomorphic to R then $S \leq Q$. Let $\psi : R \rightarrow S$ be an isomorphism. Then $\psi\Phi$ is an isomorphism in \mathcal{E} , and since $S\Phi$ is trivial, we must have that $R\Phi$ is trivial also. Thus Q is strongly \mathcal{F} -closed. \square

We will now construct, for every strongly \mathcal{F} -closed subgroup Q , a morphism on \mathcal{F} with kernel exactly Q .

Definition 5.6 Let P be a finite p -group and let \mathcal{F} be a fusion system on P . Let Q be a strongly \mathcal{F} -closed subgroup of P . By $\bar{\mathcal{F}}_Q$, we will denote the subobject of the universal fusion system $\mathcal{U}(P/Q)$, containing all objects in $\mathcal{U}(P/Q)$, and with morphisms $\text{Hom}_{\bar{\mathcal{F}}_Q}(R/Q, S/Q)$ consisting of those morphisms induced by $\text{Hom}_{\mathcal{F}}(R', S')$, as R' and S' range over all subgroups of P such that $R'Q = R$ and $S'Q = S$. (Since Q is strongly \mathcal{F} -closed, any such morphism $\phi : R' \rightarrow S'$ gives rise to a morphism $\bar{\phi} : R'Q/Q \rightarrow S'Q/Q$ using $R'Q/Q \cong R'/R' \cap Q$.)

By $\langle \bar{\mathcal{F}}_Q \rangle$ we denote the category on P/Q consisting of all compositions of morphisms from $\bar{\mathcal{F}}_Q$.

Notice that \mathcal{F}/Q is contained inside $\bar{\mathcal{F}}_Q$. It turns out that $\langle \bar{\mathcal{F}}_Q \rangle$ is a fusion system.

Lemma 5.7 Let \mathcal{F} be a fusion system on a finite p -group P . If Q is a strongly \mathcal{F} -closed subgroup of P , then $\langle \bar{\mathcal{F}}_Q \rangle$ is a fusion system on P/Q .

Proof: That $\langle \bar{\mathcal{F}}_Q \rangle$ is a category is obvious, since we are guaranteed compositions of morphisms by definition. The first axiom of a fusion system is satisfied, since

$$\mathcal{F}_{P/Q}(P/Q) \subseteq \mathcal{F}/Q \subseteq \langle \bar{\mathcal{F}}_Q \rangle.$$

If we prove the final two axioms for the subset $\bar{\mathcal{F}}_Q$, then since $\langle \bar{\mathcal{F}}_Q \rangle$ is got from $\bar{\mathcal{F}}_Q$ by compositions of morphisms, those axioms would also hold there. If $\phi : R/Q \rightarrow S/Q$ is a morphism in $\bar{\mathcal{F}}_Q$, then there is some morphism $\phi' : R' \rightarrow S'$ inducing ϕ , and the corresponding isomorphism $\psi' : R' \rightarrow R'\phi'$ induces an isomorphism in $\bar{\mathcal{F}}_Q$ corresponding to ϕ . Finally, if $\phi : R/Q \rightarrow S/Q$ is an isomorphism in $\bar{\mathcal{F}}_Q$, then it comes from some isomorphism $\phi' : R' \rightarrow S'$ in \mathcal{F} , and the inverse of ϕ' induces the inverse of ϕ . Hence $\langle \bar{\mathcal{F}}_Q \rangle$ is a fusion system. \square

At this point it becomes difficult to know whether to define the natural map $\mathcal{F} \rightarrow \langle \bar{\mathcal{F}}_Q \rangle$ as a morphism of fusion systems, since although it satisfies the requirements in the definition, the image of the map, $\bar{\mathcal{F}}_Q$, is not in general a fusion system, because it is not a category. If $\bar{\mathcal{F}}_Q = \langle \bar{\mathcal{F}}_Q \rangle$, then the natural map does become a surjective morphism of fusion systems.

Proposition 5.8 Let \mathcal{F} be a fusion system on a finite p -group P , and suppose that Q is a subgroup such that $\mathcal{F} = N_{\mathcal{F}}(Q)$. Then $\mathcal{F}/Q = \bar{\mathcal{F}}_Q$, and hence the natural map $\mathcal{F} \rightarrow \mathcal{F}/Q$ is a morphism of fusion systems.

Proof: Any morphism $\phi : R \rightarrow S$ extends to a morphism $\psi : QR \rightarrow QS$ that acts as an automorphism on Q . Certainly, $\bar{\phi} = \bar{\psi}$ in $\bar{\mathcal{F}}_Q$, since the action of ϕ and ψ on QR/Q is the same. Also, $\bar{\psi} \in \mathcal{F}/Q$, and since $\mathcal{F}/Q \subseteq \bar{\mathcal{F}}_Q$ we must have equality. \square

In general of course, a strongly \mathcal{F} -closed subgroup need not be normal in \mathcal{F} , and in this case we need not have that $\bar{\mathcal{F}}_Q$ is a fusion system, or that $\langle \bar{\mathcal{F}}_Q \rangle = \mathcal{F}/Q$.

Example 5.9 Let $P = \langle a, b, c, d \rangle$ be elementary abelian of order 16, and let \mathcal{F} be the fusion system generated by $\mathcal{F}_P(P)$ and the two morphisms $\langle ab \rangle \rightarrow \langle c \rangle$ and $\langle ac \rangle \rightarrow \langle d \rangle$. Then $A = \langle a \rangle$ is strongly \mathcal{F} -closed, and so we may form the object $\bar{\mathcal{F}}_A$. Here, the cosets Ab and Ac are $\bar{\mathcal{F}}_A$ -conjugate, as are the cosets Ac and Ad . However, there is no map sending Ab to Ad , and so $\bar{\mathcal{F}}_A$ is not a fusion system.

Note also that there are no non-trivial morphisms on overgroups of A , and so $\mathcal{F}/A = \mathcal{F}_{P/A}(P/A)$.

Other than the case where $\mathcal{F} = N_{\mathcal{F}}(Q)$, there is another case in which $\mathcal{F}/Q = \bar{\mathcal{F}}_Q$. The proof of this theorem follows [12, Proposition 6.3], although simplifications have been made.

Theorem 5.10 Let \mathcal{F} be a saturated fusion system and let Q be a strongly \mathcal{F} -closed subgroup. Then $\mathcal{F}/Q = \bar{\mathcal{F}}_Q$, and so the map $\mathcal{F} \rightarrow \mathcal{F}/Q$ is a morphism of fusion systems.

Proof: If $\phi : R \rightarrow S$ is a map in \mathcal{F} , write $\bar{\phi}$ for the image of this map in $\bar{\mathcal{F}}_Q$; i.e., write $\bar{\phi}$ for the induced map $\bar{\phi} : QR/Q \rightarrow QS/Q$. Firstly, notice that both \mathcal{F}/Q and $\bar{\mathcal{F}}_Q$ are defined on the same group, namely P/Q . Certainly, \mathcal{F}/Q is contained in $\bar{\mathcal{F}}_Q$, so we need to prove the converse; in other words, given a morphism $\phi : R \rightarrow S$ in \mathcal{F} , we need to show that there is some morphism $\psi : RQ \rightarrow P$ such that $\bar{\phi} = \bar{\psi}$, for then $\bar{\phi} \in \mathcal{F}/Q$, as needed.

We proceed by induction on $n = |P : R|$, noting that if R contains Q then we are done trivially; in particular, this implies that $n > 1$. By Alperin's fusion theorem, any morphism may be factored as (restrictions of) a sequence of automorphisms ϕ_i of fully normalized, \mathcal{F} -centric, \mathcal{F} -radical subgroups U_i . Suppose that the images $\bar{\phi}_i$ of each of the ϕ_i lie in \mathcal{F}/Q : since \mathcal{F}/Q is a fusion system and $\mathcal{F} \rightarrow \langle \bar{\mathcal{F}}_Q \rangle$ is a morphism, we have that $\bar{\phi}$, the product of (restrictions of) the $\bar{\phi}_i$, also lies in \mathcal{F}/Q .

Therefore we may assume that one of the $\bar{\phi}_i$ lies in $\bar{\mathcal{F}}_Q$ but not in \mathcal{F}/Q . By our inductive hypothesis, we see that $|U_i| = |R|$, and so we may replace R and ϕ by U_i and ϕ_i ; therefore R is now a fully normalized, \mathcal{F} -centric, \mathcal{F} -radical subgroup, and ϕ is an automorphism of R . Also, $Q \not\leq R$ as we saw above.

Similar to the proof of Proposition 5.3, let K be the kernel of the natural map $\text{Aut}_{\mathcal{F}}(R) \rightarrow \text{Aut}(RQ/Q)$, a normal subgroup of $A = \text{Aut}_{\mathcal{F}}(R)$; then K consists of all elements of $\text{Aut}_{\mathcal{F}}(R)$ that

act trivially on QR/Q , and hence are sent to the identity automorphism of RQ/Q under the map $\mathcal{F} \rightarrow \mathcal{F}/Q$.

Let $T = N_P^K(R)$; since R is fully normalized, $\text{Aut}_P(R)$ is a Sylow p -subgroup of A , and so $K \cap \text{Aut}_P(R) = \text{Aut}_T(R)$ is a Sylow p -subgroup of K . Therefore, by the Frattini argument,

$$A = K N_A(\text{Aut}_T(R)).$$

Step 1: We have $Q \cap T = N_Q^K(R) = N_Q(R)$, and $RN_Q(R) > R$. The first equality is obvious. Let $g \in N_Q(R)$, and $x \in R$. Then it is easy to see that $Qx^g = Qx$, so that g acts trivially on QR/Q . Hence the automorphism determined by g is in K , and so our first claim is proved. To see the second part, notice that $Q \cap R < Q$, and so $N_Q(R) = N_Q(Q \cap R) > Q \cap R$.

Step 2: If $\psi \in N_A(\text{Aut}_T(R))$, then N_ψ contains T . Since N_ψ is the inverse image under $\delta_R : N_P(R) \rightarrow \text{Aut}(R)$ of the subgroup $\text{Aut}_P(R) \cap \text{Aut}_P(R)^{\psi^{-1}}$ in A , we need to show that $\text{Aut}_T(R)$ is contained in both terms of the intersection. That it is contained in the first is clear, and for the second, since $\psi \in N_A(\text{Aut}_T(R))$, we have that $(\text{Aut}_T(R))^\psi = \text{Aut}_T(R)$. Thus it is contained in both terms, and so our claim is proved.

Now we may prove the result: since $A = K N_A(\text{Aut}_T(R))$, the morphism ϕ may be written as $\phi = \chi\psi$, where $\chi \in K$ and $\psi \in N_A(\text{Aut}_T(R))$. Since χ acts trivially on QR/Q , we see that $\bar{\phi} = \bar{\psi}$ in $\bar{\mathcal{F}}_Q$. Furthermore, by Step 2 we see that N_ψ contains T . However, if $N_\psi > R$, then ψ extends to ψ' , on an overgroup of R , and so by induction $\bar{\psi}'$ lies in \mathcal{F}/Q . Since \mathcal{F}/Q is a fusion system, this would imply that $\bar{\psi}$ is in \mathcal{F}/Q . Therefore $T \leq R$, and hence $Q \cap T = N_Q(R) \leq R$. However, by Step 1, $RN_Q(R) > R$, a contradiction, proving the theorem. \square

Thus if \mathcal{F} is a saturated fusion system and Q is a strongly \mathcal{F} -closed subgroup, then one may use either of the systems \mathcal{F}/Q or $\bar{\mathcal{F}}_Q$ when making arguments about quotient systems. This will be essential in our short proof of Theorem D.

At this point we need to make a remark about the factor system \mathcal{F}/Q ; Markus Linckelmann, in private communication, has pointed out that $\mathcal{F}/Q = N_{\mathcal{F}}(Q)/Q$, and so \mathcal{F}/Q is determined *locally*. The interaction between this statement and Theorem 5.10 might have significant implications for the structure of fusion systems, which have not yet been considered.

We end this section with what are essentially the second and third isomorphism theorems for fusion systems. Recall the definition of the universal fusion system on a finite p -group, consisting of all injective homomorphisms between subgroups of P .

Proposition 5.11 (Second Isomorphism Theorem) Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a strongly \mathcal{F} -closed subgroup, and let \mathcal{E} be a saturated subsystem on a subgroup R on P . Write $\mathcal{E}Q/Q$ for the image of \mathcal{E} in $\bar{\mathcal{F}}_Q$. Then

$$\mathcal{E}Q/Q \cong \mathcal{E}/(R \cap Q).$$

Proof: The isomorphism $RQ/Q \rightarrow R/R \cap Q$ induces an isomorphism $\Phi : \mathcal{U}(RQ/Q) \rightarrow \mathcal{U}(R/R \cap Q)$ of the universal fusion systems, so we need to prove that the image of $\mathcal{E}Q/Q$ in $\mathcal{U}(R/R \cap Q)$ lies inside $\mathcal{E}/R \cap Q$ and vice versa.

Let S/Q and T/Q be subgroups of RQ/Q . Then a morphism $\phi : S/Q \rightarrow T/Q$ lies in $\mathcal{E}Q/Q$ if and only if there exist subgroups S' and T' of R with $S'Q = S$ and $T'Q = T$, and a morphism $\psi \in \text{Hom}_{\mathcal{E}}(S', T')$ such that the image of ψ in $\bar{\mathcal{F}}_Q$ is ϕ . The image of ψ in $\mathcal{E}/R \cap Q$ is clearly $\phi\Phi$, and so the image of $\mathcal{E}Q/Q$ under Φ is contained in $\mathcal{E}/R \cap Q$. Conversely, if $\theta : S/R \cap Q \rightarrow T/R \cap Q$ is a morphism in $\mathcal{E}/R \cap Q$, then there is a morphism $\chi : S \rightarrow T$ in \mathcal{E} with image θ in $\mathcal{E}/R \cap Q$, and the image $\bar{\chi}$ of χ in $\mathcal{E}Q/Q$ also satisfies $\bar{\chi}\Phi = \theta$, and so Φ induces an isomorphism $\mathcal{E}Q/Q \rightarrow \mathcal{E}/R \cap Q$, as needed. \square

We get the following corollary *a posteriori*.

Corollary 5.12 Let \mathcal{F} be a saturated fusion system, and let Q be a strongly \mathcal{F} -closed subgroup. The image of any saturated subsystem of \mathcal{F} in $\bar{\mathcal{F}}_Q$ is saturated.

We now consider the third isomorphism theorem for fusion systems.

Proposition 5.13 (Third Isomorphism Theorem) Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q and R be strongly \mathcal{F} -closed subgroups with $Q \leq R$. Then

$$(\mathcal{F}/Q)/(R/Q) \cong \mathcal{F}/R.$$

Proof: By the third isomorphism theorem for groups, the two fusion systems $\mathcal{E} = (\mathcal{F}/Q)/(R/Q)$ and \mathcal{F}/R are on the same subgroup. Suppose that $\bar{\phi} : S/R \rightarrow T/R$ is a morphism in \mathcal{F}/R . Then there is some morphism $\phi \in \text{Hom}_{\mathcal{F}}(S, T)$ with image $\bar{\phi}$. Furthermore, the image $\phi' : S/Q \rightarrow T/Q$ of ϕ in \mathcal{F}/Q has image $\phi'' : S/R \rightarrow T/R$ in $(\mathcal{F}/Q)/(R/Q)$, and since both $\bar{\phi}$ and ϕ'' are derived from ϕ , they must be the same morphism. The converse is a similar calculation, and is safely omitted. \square

In the second isomorphism theorem, we denoted by $\mathcal{E}Q/Q$ the image of \mathcal{E} in \mathcal{F}/Q . If \mathcal{E} is defined on a subgroup R of P , then there is a subsystem $\mathcal{E}Q$, defined on RQ , such that the factor system $\mathcal{E}Q/Q$ is the image of \mathcal{E} in \mathcal{F}/Q , namely the full preimage under the natural morphism $\mathcal{F} \rightarrow \mathcal{F}/Q$. It is not clear at this time whether there is always a *saturated* such subsystem $\mathcal{E}Q$, but it is known that fusion systems do not behave as well as groups with respect to taking preimages, so it seems unlikely that this is always true. (In the case where $\mathcal{F} = \text{C}_{\mathcal{F}}(Q)$, there is a bijection between saturated subsystems of \mathcal{F} on subgroups containing Q and saturated subsystems of \mathcal{F}/Q : see [6, Section 6].)

6 Closure and Quotients

We begin with how weak and strong closure relates to taking quotients. The proof in [13] of Theorem 6.1(iv) given below is not clear, because it is only true given Theorem 5.10, a result that is not mentioned in [13].

Theorem 6.1 (cf. Stancu [13, Lemma 4.7], Aschbacher [4, Lemma 8.9]) Let \mathcal{F} be a fusion system on a finite p -group P , and let Q be a strongly \mathcal{F} -closed subgroup of P , and let R be a subgroup of P .

- (i) The map $\Phi : \mathcal{F} \rightarrow \mathcal{F}/Q$ induces a bijection between the weakly \mathcal{F} -closed subgroups of P containing Q and the weakly \mathcal{F}/Q -closed subgroups of P/Q .
- (ii) If R is weakly \mathcal{F} -closed then the image of R in \mathcal{F}/Q is weakly \mathcal{F}/Q -closed.
- (iii) If \mathcal{F} is saturated then the map $\Phi : \mathcal{F} \rightarrow \mathcal{F}/Q$ induces a bijection between the strongly \mathcal{F} -closed subgroups of P containing Q and the strongly \mathcal{F}/Q -closed subgroups of P/Q .
- (iv) If \mathcal{F} is saturated and R is strongly \mathcal{F} -closed then the image of R in \mathcal{F}/Q is strongly \mathcal{F}/Q -closed.

Proof: Let R be a subgroup of P containing Q . Then R is weakly closed if and only if any morphism $\phi : R \rightarrow P$ is an automorphism. Clearly if R is weakly \mathcal{F} -closed then any morphism $\phi' : R/Q \rightarrow P/Q$ in \mathcal{F}/Q (which must have a preimage in $\text{Hom}_{\mathcal{F}}(R, P)$) is an automorphism, and vice versa. Hence (i) is true.

Also, the product of two weakly \mathcal{F} -closed subgroups is weakly \mathcal{F} -closed, so if any subgroup R is weakly \mathcal{F} -closed, so is QR , and hence QR/Q is weakly \mathcal{F}/Q -closed by (i), proving (ii).

For the rest of the proof, suppose that \mathcal{F} is saturated, so that $\bar{\mathcal{F}}_Q = \mathcal{F}/Q$. Let R be an overgroup of Q , and let S be any subgroup of R . Suppose that $\phi : S \rightarrow P$ is a morphism in \mathcal{F} . Then $S\phi \leq R$ if and only if $(SQ/Q)\bar{\phi} = (S\phi)Q/Q \leq R/Q$, where $\bar{\phi}$ is the image in $\bar{\mathcal{F}}_Q$ of ϕ . Since this holds for all ϕ , we see that R is strongly \mathcal{F} -closed if and only if R/Q is strongly $(\bar{\mathcal{F}}_Q = \mathcal{F}/Q)$ -closed, proving (iii).

Let R be any strongly \mathcal{F} -closed subgroup of P , let S be any subgroup of R , and let $\phi \in \text{Hom}_{\mathcal{F}}(SQ, P)$. Since Q is strongly \mathcal{F} -closed, $\phi|_Q$ is an automorphism of Q , and since R is strongly \mathcal{F} -closed, $\phi|_S$ maps S to R . Therefore, $(SQ)\phi \leq RQ$. If every subgroup of RQ/Q is of the form SQ/Q for some $S \leq R$, then RQ/Q would be strongly \mathcal{F}/Q -closed, proving (iv). However, this follows from the second isomorphism theorem, since $RQ/Q \cong R/R \cap Q$, choosing A/Q corresponds to a subgroup $B/R \cap Q$ on the right-hand side, and $BQ/Q = A/Q$, as needed. \square

We now prove Theorem C, by collating the equivalent conditions to a subgroup Q having the property that $\mathcal{F} = N_{\mathcal{F}}(Q)$.

Theorem 6.2 Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a subgroup of P . The following are equivalent:

- (i) $\mathcal{F} = N_{\mathcal{F}}(Q)$;
- (ii) $\mathcal{F}_Q(Q) \prec \mathcal{F}$;
- (iii) Q is contained in every fully normalized, \mathcal{F} -centric, \mathcal{F} -radical subgroup of \mathcal{F} ;
- (iv) there is a central series for Q all of whose terms (including Q) are strongly \mathcal{F} -closed; and
- (v) there is a central series for Q all of whose terms are weakly \mathcal{F} -closed, and Q is strongly \mathcal{F} -closed.

Proof: The equivalence of (i) and (ii) is Theorem A, the equivalence of (i) and (iii) is Proposition 3.3, the equivalence of (i) and (iv) is Proposition 3.4, and that (iv) implies (v) is obvious. It remains to show that (v) implies (iv).

Let

$$1 = Q_0 \leq Q_1 \leq \cdots \leq Q_n = Q$$

be a central series for Q , all of whose terms are weakly \mathcal{F} -closed. By Lemma 3.7, $Q_1 \leq Z(Q)$ is strongly \mathcal{F} -closed. Since Q_1 is strongly \mathcal{F} -closed, we may take the quotient system \mathcal{F}/Q_1 . Theorem 6.1 states that the map $\mathcal{F} \rightarrow \mathcal{F}/Q_1$ induces a bijection between the weakly and strongly \mathcal{F} -closed subgroups of P containing Q_1 and the weakly and strongly \mathcal{F}/Q_1 -closed subgroups of P/Q_1 respectively.

At this stage, one may either proceed by induction on the length of a central series all of whose terms are weakly \mathcal{F} -closed, by noticing that now Q_2/Q_1 is strongly \mathcal{F}/Q_1 -closed, and hence Q_2 is strongly \mathcal{F} -closed, or proceed by induction on $|Q|$, and note that since Q/Q_1 satisfies (iv) of the theorem, there must be a central series by (v) which, when full preimages are taken, gives a central series for Q all of whose terms are strongly \mathcal{F} -closed. Hence all conditions are equivalent, as claimed. \square

Using Theorem 6.1, the proof of Theorem D is trivial. Let Q and R be strongly \mathcal{F} -closed subgroups of a saturated fusion system \mathcal{F} . By Theorem 6.1(iv), QR/Q is strongly \mathcal{F}/Q -closed, and by (iii) of that theorem, this implies that QR is strongly \mathcal{F} -closed, as required.

We now include a couple of lemmas, ready for our treatment of p -soluble fusion systems. If \mathcal{F} is a fusion system on P , then the *centre* of \mathcal{F} , denoted $Z(\mathcal{F})$, is the collection of all elements $x \in P$ such that $C_{\mathcal{F}}(\langle x \rangle) = \mathcal{F}$.

Lemma 6.3 Let \mathcal{F} be a saturated fusion system on a finite p -group P , and suppose that Z is a strongly \mathcal{F} -closed subgroup of $Z(\mathcal{F})$. Then for any normal subgroup Q containing Z , $\mathcal{F} = N_{\mathcal{F}}(Q)$ if and only if $\mathcal{F}/Z = N_{\mathcal{F}/Z}(Q/Z)$.

Proof: This is a special case of [11, Theorem 6.5]. \square

Lemma 6.4 Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a fully normalized subgroup. If K is a normal subgroup of $\text{Aut}_{\mathcal{F}}(Q)$; then $N_{\mathcal{F}}^K(Q) \triangleleft N_{\mathcal{F}}(Q)$. In particular, $C_{\mathcal{F}}(Q) \triangleleft N_{\mathcal{F}}(Q)$.

Proof: Since \mathcal{F} is saturated and Q is fully normalized, $N_{\mathcal{F}}(Q)$ is saturated. Also, since K is a normal subgroup of $\text{Aut}_{\mathcal{F}}(Q)$, $N_{\mathcal{F}}^K(Q)$ is saturated as well, as we remarked when we defined $N_{\mathcal{F}}^K(Q)$.

Next, we need to show that $N_{\mathcal{F}}^K(Q)$ is strongly $N_{\mathcal{F}}(Q)$ -closed, so let R be a subgroup of $N = N_{\mathcal{F}}^K(Q)$ and $\phi : R \rightarrow S$ be a morphism in $N_{\mathcal{F}}(Q)$. For $g \in R$, consider the conjugation action c_g of g on Q ; since $R \leq N$, we see that $c_g \in K$. Furthermore, the action of $g\phi \in S$ on Q is given by

$$c_{g\phi} = \phi^{-1}c_g\phi \in K,$$

since $K \triangleleft \text{Aut}_{\mathcal{F}}(Q)$. Therefore $S \leq N_{\mathcal{F}}^K(Q)$, and so N is strongly \mathcal{F} -closed.

Finally, we need to show that $N_{\mathcal{F}}^K(Q)$ is $N_{\mathcal{F}}(Q)$ -invariant, so let $R \leq S \leq N$, $\phi \in \text{Hom}_{N_{\mathcal{F}}^K(Q)}(R, S)$ and $\psi \in \text{Hom}_{N_{\mathcal{F}}(Q)}(S, N)$. Since each of ϕ and ψ extends to maps $\bar{\phi}$ and $\bar{\psi}$ whose domains include Q , and in the first case $\bar{\phi}|_Q \in K$ and in the second $\bar{\psi}|_Q \in \text{Aut}_{\mathcal{F}}(Q)$, we see that

$$(\bar{\psi}^{-1}\bar{\phi}\bar{\psi})|_Q \in K,$$

as K is a normal subgroup of $\text{Aut}_{\mathcal{F}}(Q)$. Hence $\psi^{-1}\phi\psi$ extends to a map θ whose domain includes Q and for which $\theta|_Q \in K$. Therefore $\psi^{-1}\phi\psi \in N_{\mathcal{F}}^K(Q)$, and so $N_{\mathcal{F}}^K(Q)$ is $N_{\mathcal{F}}(Q)$ -invariant, as required. \square

7 p -Soluble Fusion Systems

For finite groups, a group is called p -soluble if repeated quotienting by (alternating) $O_p(G)$ and $O_{p'}(G)$ reaches the identity. In the case of fusion systems, we have that $O_{p'}(\mathcal{F}) = 1$, so it makes sense to make the following definition.

Definition 7.1 Let \mathcal{F} be a saturated fusion system on a finite p -group P . We say that \mathcal{F} is p -soluble if there exists a chain of strongly \mathcal{F} -closed subgroups

$$1 = P_0 \leq P_1 \leq \dots \leq P_n = P,$$

such that $P_i/P_{i-1} \leq O_p(\mathcal{F}/P_{i-1})$ for all $1 \leq i \leq n$. If \mathcal{F} is p -soluble, then the length n of a smallest such chain above will be called the p -length of \mathcal{F} .

We have the following easy lemma.

Lemma 7.2 Let \mathcal{F} be a saturated fusion system on a finite p -group P . If Q is a strongly \mathcal{F} -closed subgroup of P , then $O_p(\mathcal{F})/Q \leq O_p(\mathcal{F}/Q)$, and if $Q \leq O_p(\mathcal{F})$, then \mathcal{F} is p -soluble if and only if \mathcal{F}/Q is p -soluble.

Proof: Let R be a strongly \mathcal{F} -closed subgroup of P with $\mathcal{F} = N_{\mathcal{F}}(R)$. We claim that RQ/Q is contained in $O_p(\mathcal{F}/Q)$. To see this, let $\psi : A/Q \rightarrow B/Q$ be any morphism in \mathcal{F}/Q , and let $\phi : A \rightarrow B$ be an \mathcal{F} -morphism whose image in \mathcal{F}/Q is ψ . This may be extended to a morphism $\bar{\phi} : AR \rightarrow BR$ with $\bar{\phi}|_R$ an automorphism; $\bar{\psi}$ of $\bar{\phi}$ is a map $AR/Q \rightarrow BR/Q$ that must act as an automorphism of RQ/Q , since $\bar{\phi}$ acts as an automorphism of R . This proves that $O_p(\mathcal{F})/Q \leq O_p(\mathcal{F}/Q)$.

For the rest of the proof, suppose that $Q \leq O_p(\mathcal{F})$. If \mathcal{F}/Q is p -soluble, then taking the preimages of a chain of strongly \mathcal{F}/Q -closed subgroups witnessing the p -solubility of \mathcal{F}/Q , together with Q itself, yields a chain of strongly \mathcal{F} -closed subgroups (by Theorem 6.1(iii)) witnessing the p -solubility of \mathcal{F} , proving one direction.

Conversely, suppose that \mathcal{F} is p -soluble, and let

$$1 = P_0 \leq P_1 \leq \cdots \leq P_n = P$$

be a sequence of strongly \mathcal{F} -closed subgroups with $P_i/P_{i-1} \leq O_p(\mathcal{F}/P_{i-1})$ for all $1 \leq i \leq n$. We may assume that P_1 is non-trivial, and notice that \mathcal{F}/P_1 is clearly p -soluble, with $|P/P_1| < |P|$. Define $\bar{Q} = P_1Q$: by induction, since $\bar{Q}/P_1 \leq O_p(\mathcal{F}/P_1)$ by the first part, and \mathcal{F}/P_1 is p -soluble, we have that $\mathcal{F}/\bar{Q} = (\mathcal{F}/P_1)/(\bar{Q}/P_1)$ is p -soluble. If $Q \neq 1$, then $|P/Q| < |P|$, so we can use induction, and therefore \mathcal{F}/Q is p -soluble if and only if $(\mathcal{F}/Q)/(\bar{Q}/Q) \cong \mathcal{F}/\bar{Q}$ is p -soluble (as $\bar{Q}/Q \leq O_p(\mathcal{F}/Q)$ by the first part). Since \mathcal{F}/\bar{Q} is p -soluble, this completes the proof. \square

The following lemma describes the basic facts of p -soluble fusion systems, mirroring those of finite groups. Define $O_p^{(0)}(\mathcal{F}) = 1$, and the i th term by

$$O_p^{(i)}(\mathcal{F})/O_p^{(i-1)}(\mathcal{F}) = O_p\left(\mathcal{F}/O_p^{(i-1)}(\mathcal{F})\right).$$

Lemma 7.3 Let \mathcal{F} be a saturated fusion system on a finite p -group P , and let Q be a strongly \mathcal{F} -closed subgroup of P .

- (i) If \mathcal{F} is p -soluble then all saturated subsystems and quotients \mathcal{F}/Q are p -soluble.
- (ii) Let \mathcal{E} be a weakly normal subsystem of \mathcal{F} , on the subgroup Q . If both \mathcal{E} and \mathcal{F}/Q are p -soluble then so is \mathcal{F} .
- (iii) \mathcal{F} is p -soluble if and only if $O_p^{(n)}(\mathcal{F}) = P$ for some n , and the smallest such n is the p -length of \mathcal{F} .

Proof: Choose a triple (\mathcal{F}, P, Q) , where \mathcal{F} is a p -soluble, saturated fusion system on a finite p -group P , and Q is a strongly \mathcal{F} -closed subgroup of P such that \mathcal{F}/Q is not p -soluble, and such that $|P|$ is minimal subject to these constraints. Write $R = O_p(\mathcal{F}) \neq 1$; by Lemma 7.2, we see that \mathcal{F}/R is p -soluble, and \mathcal{F}/QR is not p -soluble (as $QR/Q \leq O_p(\mathcal{F}/Q)$ and hence \mathcal{F}/Q is p -soluble if and only if $(\mathcal{F}/Q)/(QR/Q) \cong \mathcal{F}/QR$ is, by another application of Lemma 7.2). This

proves that $(\mathcal{F}/R, P/R, QR/R)$ is a triple satisfying our conditions with $|P| > |P/R|$. This yields a contradiction, proving that quotients of p -soluble fusion systems are p -soluble.

Now let $(\mathcal{F}, \mathcal{F}', Q)$ be a triple, with \mathcal{F} a p -soluble fusion system on a p -group P , \mathcal{F}' a saturated subsystem of \mathcal{F} on a subgroup Q , with \mathcal{F}' not p -soluble. Choose this triple with $|P|$ minimal. Let $R = O_p(\mathcal{F}) \neq 1$, and we claim that $R \cap Q \leq O_p(\mathcal{F}')$. Since $N_{\mathcal{F}}(R) = \mathcal{F}$, we have a central series

$$1 = R_0 \leq R_1 \leq \cdots \leq R_n = R,$$

with each R_i strongly \mathcal{F} -closed, by Proposition 3.4. Therefore any morphism in \mathcal{F}' whose domain lies inside R_i has image inside R_i . Since Q is obviously strongly \mathcal{F}' -closed, this means that the intersection $Q_i = R_i \cap Q$ is strongly \mathcal{F}' -closed. Thus the series of the Q_i is a central series for $Q \cap R$ whose terms are strongly \mathcal{F}' -closed, which by another application of Proposition 3.4, gives the result.

Consider the image $\mathcal{F}'R/R$ of \mathcal{F}' in \mathcal{F}/R . Since this is isomorphic with $\mathcal{F}'/R \cap Q$ by the second isomorphism theorem, and is hence not p -soluble (by Lemma 7.2 again), the triple $(\mathcal{F}/R, \mathcal{F}'R/R, QR/R)$ has $|P/R| < |P|$ and so this contradicts the choice of the original triple. This proves (i).

Recall that by $O_p(\mathcal{E})$ we mean the subsystem $\mathcal{F}_{O_p(\mathcal{E})}(O_p(\mathcal{E}))$. We have that $O_p(\mathcal{E}) \text{ char } \mathcal{E} \prec \mathcal{F}$, so that $O_p(\mathcal{E})$ is strongly \mathcal{F} -closed; this, together with induction and Theorem 6.1(iii), proves that $O_p^{(i)}(\mathcal{E})$ is strongly \mathcal{F} -closed for all i . Since \mathcal{E} is p -soluble $O_p^{(i)}(\mathcal{E})/O_p^{(i-1)}(\mathcal{E}) \leq O_p(\mathcal{E}/O_p^{(i-1)}(\mathcal{E}))$, and Lemma 4.1 therefore implies that $O_p^{(i)}(\mathcal{E})/O_p^{(i-1)}(\mathcal{E}) \leq O_p(\mathcal{F}/O_p^{(i-1)}(\mathcal{E}))$; this chain of strongly \mathcal{F} -closed subgroups satisfies the requirements for \mathcal{F} to be p -soluble up to Q , the subgroup on which \mathcal{E} lies. By Theorem 6.1(iii), the preimages of $O_p^{(j)}(\mathcal{F}/Q)$ are strongly \mathcal{F} -closed. The concatenation of the two series $O_p^{(i)}(\mathcal{E})$ and the preimages of $O_p^{(j)}(\mathcal{F}/Q)$ satisfy the requirement for \mathcal{F} to be p -soluble, proving (ii).

Now, we prove (iii). One direction is clear, so suppose that \mathcal{F} has a series (Q_i) so that $Q_i/Q_{i-1} \leq O_p(\mathcal{F}/Q_{i-1})$ for all i . If we can show that, whenever $S \leq O_p(\mathcal{F})$ is strongly \mathcal{F} -closed, then the preimage of $O_p(\mathcal{F}/S)$ is contained in $O_p^{(2)}(\mathcal{F})$, then we are done by an obvious induction.

By Lemma 7.2, if $S \leq O_p(\mathcal{F}) = R$, then $R/S \leq O_p(\mathcal{F}/S)$, and so one may consider $O_p(\mathcal{F}/S)/(R/S)$, which is a subgroup of $O_p((\mathcal{F}/S)/(R/S)) \cong O_p(\mathcal{F}/R)$ by another application of Lemma 7.2, and hence contained in $O_p^{(2)}(\mathcal{F})/R$. Thus all parts of the lemma are proved. \square

At this point we digress briefly to discuss the concept of minimal weakly normal subsystems, i.e., weakly normal subsystems of a fusion system \mathcal{F} that contain no other weakly normal subsystem apart from themselves and 1. At the moment, there is no characterization of minimal weakly normal subsystems, like there is for groups. The reason behind this is that the intersection of two weakly normal subsystems need not be weakly normal, nor even saturated.

Example 7.4 Let $P = D_8 \times C_2$, with the D_8 factor generated by an element x of order 4 and y of order 2, and the C_2 -factor being generated by z . Let $Q = \langle x, y \rangle$, and $R = \langle xz, y \rangle$. Then $S = Q \cap R$ is a normal Klein four subgroup of P , and $\text{Aut}_Q(S) = \text{Aut}_R(S)$ contains the identity and the map

swapping y and x^2y . Thus $\mathcal{E} = \mathcal{F}_Q(Q) \cap \mathcal{F}_R(R)$ has an outer automorphism of order 2 on S , and so cannot be saturated, as $\text{Aut}_S(S)$ is not a Sylow 2-subgroup of $\text{Aut}_{\mathcal{E}}(S)$.

This example shows that for any ‘reasonable’ definition of normality – i.e., one for which normal subgroups yield normal subsystems – the intersection of two normal subsystems need not be saturated. As we mentioned in Section 2, in [3] Aschbacher proves that the intersection of two normal subsystems *contains* a normal subsystem on the intersection of the two relevant subgroups. This result easily enables one to prove (mimicking the proof for groups) that a minimal normal subsystem of a saturated fusion system is either the fusion system $\mathcal{F}_Q(Q)$ for an elementary abelian p -subgroup Q or a direct product of isomorphic *non-abelian* simple subsystems, that is, simple subsystems not of the form $\mathcal{F}_R(R)$ for $R \cong C_p$. Such a result is not known for weakly normal subsystems.

However, we can say something about minimal *weakly subnormal* subsystems. (The definition of weak subnormality is obvious, and left to the reader.) These are obviously simple, and so either of the form $\mathcal{F}_Q(Q)$ for Q of prime order, or some non-abelian simple fusion system.

Proposition 7.5 Let \mathcal{E} be a minimal weakly subnormal subsystem of the saturated fusion system \mathcal{F} , on a finite p -group P . If $\mathcal{E} = \mathcal{F}_Q(Q)$ for $Q \leq P$ of prime order, then $Q \leq O_p(\mathcal{F})$. In particular, \mathcal{F} is not p -soluble if and only if there is a strongly \mathcal{F} -closed subgroup R and non-abelian simple subsystem of \mathcal{F}/R .

Proof: The first statement, that $O_p(\mathcal{E}) \leq O_p(\mathcal{F})$ for any weakly subnormal subsystem \mathcal{E} of \mathcal{F} , easily follows from Proposition 4.1 and induction on the length of a chain of weakly normal subsystems connecting \mathcal{E} and \mathcal{F} .

To see the second, if \mathcal{F} is p -soluble then all quotients and saturated subsystems are p -soluble, and so \mathcal{F} contains no non-abelian simple subquotient, proving one direction. For the other, proceed by induction on $|P|$; assume that \mathcal{F} is a saturated fusion system on a finite p -group P with no non-abelian simple subquotients. If $O_p(\mathcal{F}) \neq 1$, then $\mathcal{F}/O_p(\mathcal{F})$ is p -soluble by induction, whence so is \mathcal{F} . However, no minimal weakly subnormal subsystem may be non-abelian simple, and so must be of the form $\mathcal{F}_Q(Q)$, and so we see that $O_p(\mathcal{F}) \neq 1$ by the first part of the proposition. \square

We come to the main result of the section, the proof that p -soluble fusion systems are constrained. We start with a simple lemma.

Lemma 7.6 For any saturated fusion system \mathcal{F} on a finite p -group P , $Z(\mathcal{F}) \leq O_p(\mathcal{F})$, and

$$O_p(\mathcal{F})/Z(\mathcal{F}) = O_p(\mathcal{F}/Z(\mathcal{F})).$$

In particular, if $O_p(\mathcal{F}) = Z(\mathcal{F})$ then either $P = O_p(\mathcal{F})$ or \mathcal{F} is not p -soluble.

Proof: Clearly $Z(\mathcal{F}) \leq O_p(\mathcal{F})$, and by Lemma 6.3,

$$O_p(\mathcal{F})/Z(\mathcal{F}) = O_p(\mathcal{F}/Z(\mathcal{F})).$$

If $O_p(\mathcal{F}) \neq P$, then $O_p(\mathcal{F}/Z(\mathcal{F}))$ is trivial. Therefore no minimal weakly subnormal subsystem \mathcal{E} of $\mathcal{F}/Z(\mathcal{F})$ is of the form $\mathcal{F}_Q(Q)$ for some Q , and so Proposition 7.5 implies that \mathcal{F} is not p -soluble, as claimed. \square

Theorem 7.7 Suppose that \mathcal{F} is p -soluble, and let $Q = O_p(\mathcal{F})$. Then $C_P(Q) = Z(Q)$, so that \mathcal{F} is constrained.

Proof: Let $Q = O_p(\mathcal{F})$, and let $\mathcal{E} = C_{\mathcal{F}}(Q)$. Since $\mathcal{F} = N_{\mathcal{F}}(Q)$, we see that $\mathcal{E} \prec \mathcal{F}$ by Lemma 6.4. Therefore, by Proposition 4.1,

$$O_p(\mathcal{E}) = O_p(\mathcal{F}) \cap C_P(Q) = Q \cap C_P(Q) = Z(Q).$$

However, since $\mathcal{E} = C_{\mathcal{F}}(Q)$, we notice that every morphism in \mathcal{E} centralizes $Z(Q) = Q \cap C_P(Q)$, so that $Z(Q) \leq Z(\mathcal{E})$. Since $Z(\mathcal{E}) \leq O_p(\mathcal{E})$ obviously, we see that

$$O_p(\mathcal{E}) = Z(\mathcal{E}).$$

As \mathcal{E} is p -soluble (since \mathcal{F} is, by Lemma 7.3), we get by Lemma 7.6 that $C_P(Q) = O_p(C_{\mathcal{F}}(Q))$. However, $O_p(\mathcal{E}) \leq O_p(\mathcal{F})$ by Proposition 4.1, so that

$$C_P(Q) = O_p(\mathcal{E}) \leq O_p(\mathcal{F}) = Q.$$

Thus Q is \mathcal{F} -centric, as required. \square

8 Soluble Systems and Soluble Groups

The fact that p -soluble fusion systems are constrained means that they are fusion systems of finite groups, by Theorem 2.5. In this section we will characterize those fusion systems that come from p -soluble groups. Since every p -soluble group has a p -soluble fusion system, it is a necessary condition that they be constrained, so we start from here.

It is well-known (see, for example, [1, 33.12]) that if G is a simple p' -group, then any quasisimple group with quotient G is also a p' -group. This gives us the following, originally proved by Hall and Higman.

Lemma 8.1 Let G be a p -soluble group with $O_{p'}(G) = 1$. Then

$$C_G(O_p(G)) \leq O_p(G).$$

Proof: A fundamental property of the generalized Fitting subgroup is that $C_G(F^*(G)) \leq F^*(G)$ [1, 31.13]. Since G is p -soluble, any subnormal quasisimple group is a p' -group, and therefore $E(G) \leq O_{p'}(G) = 1$. Therefore $F^*(G) = O_p(G)$, proving the claim. \square

Using this lemma, we give our first characterization of fusion systems of p -soluble groups. In the rest of this section, for a constrained fusion system \mathcal{F} , we denote the unique finite group given in Theorem 2.5 by $L_{\mathcal{F}}$.

Proposition 8.2 Let \mathcal{F} be a saturated fusion system on a finite p -group P . There is a p -soluble group G such that $\mathcal{F} = \mathcal{F}_P(G)$ if and only if \mathcal{F} is constrained, and $L_{\mathcal{F}}$ is p -soluble.

Proof: Suppose that $\mathcal{F} = \mathcal{F}_P(G)$ for some p -soluble group G . We may assume that $O_{p'}(G) = 1$, since the fusion systems on G and $G/O_{p'}(G)$ are the same. By Lemma 8.1, $O_p(G)$ contains its centralizer. Therefore, by the uniqueness of the group in Theorem 2.5, $G = L_{\mathcal{F}}$. \square

Finally, this allows us to reach an internal characterization of fusion systems of p -soluble groups, without reference to groups.

Theorem G Let \mathcal{F} be a saturated fusion system on a finite p -group P , and write $Q = O_p(\mathcal{F})$. Then \mathcal{F} is the fusion system of a p -soluble group if and only if \mathcal{F} is constrained and $\text{Aut}_{\mathcal{F}}(Q)$ is p -soluble.

Proof: Suppose that \mathcal{F} is constrained and that $\text{Aut}_{\mathcal{F}}(Q)$ is p -soluble. Since \mathcal{F} is constrained, we have that $\mathcal{F} = \mathcal{F}_P(L_{\mathcal{F}})$; then $L_{\mathcal{F}}$ is an extension of $C_{L_{\mathcal{F}}}(Q) = Z(Q)$ by $\text{Aut}_{L_{\mathcal{F}}}(Q) = \text{Aut}_{\mathcal{F}}(Q)$, which is p -soluble. Hence $L_{\mathcal{F}}$ is p -soluble, and so \mathcal{F} is the fusion system of a p -soluble group. The converse is similarly clear. \square

At first blush, this appears to contradict an assertion in [12, 1.6], which claims that all p -soluble fusion systems arise from p -soluble groups. The incongruity stems from the definition of a soluble fusion system in [12], which works ‘from the top down’, in the sense that it involves taking repeated subsystems, rather than repeated quotients. Recall the definitions of $O^p(\mathcal{F})$ and $O^{p'}(\mathcal{F})$ from [12, 7.5, 6.11] (where they are denoted \mathcal{F}^{\flat} and \mathcal{F}^{α} respectively), which will not be repeated here. Then another way to define a soluble fusion system is that repeated taking of O^p and $O^{p'}$ operators eventually reaches the trivial group. While this definition picks out exactly the fusion systems of p -soluble groups (this is not difficult to prove given Theorem G), it suffers from the fact that this class of fusion system is not closed under extensions. The class of p -soluble fusion systems given here is extension-closed and contains the fusion system $\mathcal{F}_P(P)$, where P is cyclic of order p ; by Proposition 7.5, it is also the class of all systems that do not contain simple subquotients. It seems to us that this is the ‘correct’ definition of solubility for fusion systems, given the use of the word ‘soluble’ throughout algebra to mean similar concepts.

We now consider so-called $Qd(p)$ -free fusion systems. Firstly, the group $Qd(p)$ is the semidirect product $(C_p \times C_p) \rtimes \text{SL}_2(p)$, with $\text{SL}_2(p)$ acting in the natural way. A finite group is called $Qd(p)$ -free if no subquotient of it is isomorphic with $Qd(p)$. As in [10], define a saturated fusion system \mathcal{F} to be $Qd(p)$ -free if, for any fully normalized, \mathcal{F} -centric subgroup Q of P , the group $L_{N_{\mathcal{F}}(Q)}$ is $Qd(p)$ -free.

Lemma 8.3 Every $Qd(p)$ -free fusion system \mathcal{F} is p -soluble.

Proof: By [10, Proposition 6.4], if \mathcal{F} is $Qd(p)$ -free then so is $\mathcal{F}/O_p(\mathcal{F})$, so it suffices to show that $O_p(\mathcal{F}) \neq 1$; this is given to us by 7.1 of the same paper. \square

Since a $Qd(p)$ -free fusion system is p -soluble it is constrained, and hence is the fusion system of some, $Qd(p)$ -free, group. Thus any theorem known to hold for $Qd(p)$ -free groups should have an analogue for $Qd(p)$ -free fusion systems. One such example is Glauberman's ZJ-Theorem: another is Thompson Factorization.

Theorem 8.4 (Thompson Factorization) Let \mathcal{F} be a $Qd(p)$ -free fusion system on a finite p -group P , where p is odd. Then

$$\mathcal{F} = N_{\mathcal{F}}(J(P)) C_{\mathcal{F}}(\Omega_1(Z(P))),$$

where $J(P)$ is the Thompson subgroup of P .

Proof: Such a decomposition holds for $Qd(p)$ -free groups (see, for example, [9, Theorem 26.9]). Let G be the group modelling \mathcal{F} given by Theorem 2.5; the subsystem \mathcal{E}_1 of \mathcal{F} induced by $H_1 = N_G(J(P))$ is $N_{\mathcal{F}}(J(P))$, and the subsystem \mathcal{E}_2 of \mathcal{F} induced by $H_2 = C_G(\Omega_1(Z(P)))$ is $C_{\mathcal{F}}(\Omega_1(Z(P)))$ by [11, Proposition 3.16]. Since every element of G may be written as h_1h_2 , with $h_i \in H_i$, every morphism in \mathcal{F} , which is induced by some element $g = h_1h_2$, may be written as $\phi_1\phi_2$, where $\phi_i \in \mathcal{E}_i$. \square

However, a naïve rewriting of Thompson Factorization for other cases fails; for example, Glauberman proved that if $p \geq 5$, then the conclusion to the above theorem holds for all p -soluble groups, with a slight modification to the centralizer term (see [9, Theorem 26.10]). This does not carry over to p -soluble fusion systems in general; for instance, the groups $Qd(p)$ themselves do not satisfy this version of Thompson Factorization.

Acknowledgments: I would like to thank Adam Glesser for being a human sounding board for many if not all of the ideas in this paper, for pointing out some errors in the original version, and for suggesting improvements to the exposition. I also had a helpful conversation with Inna Korchagina on 2-soluble fusion systems, in which we proved Proposition 8.2 for this case. I wish to thank the referee for many helpful comments that greatly improved the exposition of the paper.

References

- [1] M. Aschbacher, *Finite Group Theory*, 2nd Edition, Cambridge University Press, Cambridge, 2000.
- [2] M. Aschbacher, *Notes on the local theory of saturated fusion systems*, from the conference 'Fusion Systems', University of Birmingham, 2007, available at <http://web.mat.bham.ac.uk/C.W.Parker/Fusion/birm0.pdf>.

- [3] M. Aschbacher, The generalized Fitting subsystem of a fusion system, preprint (2007).
- [4] M. Aschbacher, Normal subsystems of fusion systems, *Proc. Lond. Math. Soc.* 97 (2008) 239–271.
- [5] C. Broto, N. Castellana, J. Grodal, R. Levi, B. Oliver, Subgroup families controlling p -local finite groups, *Proc. London Math. Soc.* 91 (2005) 325–354.
- [6] C. Broto, N. Castellana, J. Grodal, R. Levi, B. Oliver, Extensions of p -local finite groups, *Trans. Amer. Math. Soc.* 359 (2007) 3791–3858.
- [7] C. Broto, R. Levi, B. Oliver, The homotopy theory of fusion systems, *J. Amer. Math. Soc.* 16 (2003) 779–856.
- [8] D. Gorenstein, *Finite Groups*, 2nd Edition, Chelsea Publishing Co., New York, 1980.
- [9] D. Gorenstein, R. Lyons, R. Solomon, *The Classification of the Finite Simple Groups, Number 2. Part I. Chapter G*, American Mathematical Society, Providence, RI, 1996.
- [10] R. Kessar, M. Linckelmann, ZJ theorems for fusion systems, *Trans. Amer. Math. Soc.* 360 (2008) 3093–3106.
- [11] M. Linckelmann, Introduction to fusion systems, in: *Group Representation Theory*, EPFL Press, Lausanne, 2007, pp. 79–113.
- [12] L. Puig, Frobenius categories, *J. Algebra* 303 (2006) 309–357.
- [13] R. Stancu, Control of fusion in fusion systems, *J. Algebra Appl.* 5 (2006) 817–837.