

On Tensor Products of Simple Modules for Simple Groups

David A. Craven

Received: 23 February 2011 / Accepted: 20 July 2011
© Springer Science+Business Media B.V. 2011

Abstract In an attempt to get some information on the multiplicative structure of the Green ring we study algebraic modules for simple groups, and associated groups such as quasisimple and almost-simple groups. We prove that, for almost all groups of Lie type in defining characteristic, the natural module is non-algebraic. For alternating and symmetric groups, we prove that the simple modules in p -blocks with defect groups of order p^2 are algebraic, for $p \leq 5$. Finally, we analyze nine sporadic groups, finding that all simple modules are algebraic for various primes and sporadic groups.

Keywords Algebraic module · Tensor product

Mathematics Subject Classification (2010) 20C20

1 Introduction

The tensor structure of the category of finite-dimensional kG -modules, where k is a field of characteristic p and G is a finite group, is a structure that remains largely shrouded in mystery. As a first approximation to understanding this structure, Alperin introduced the notion of *algebraic* modules [1]; a kG -module is *algebraic* if it satisfies a polynomial with integer coefficients, where addition and multiplication are given by the direct sum and tensor product. Trivial source modules are algebraic, as are simple modules for p -soluble groups [17], simple modules in characteristic 2 for finite groups with abelian Sylow 2-subgroups [8], and simple modules for 2-blocks with Klein-four defect groups in general [11].

Presented by Jon Carlson.

D. A. Craven (✉)
Mathematical Institute, University of Oxford, 24–29 St Giles', Oxford, OX1 3LB, UK
e-mail: craven@maths.ox.ac.uk

Since simple modules for p -soluble groups are algebraic, it is natural to ask about simple modules for simple groups, or groups that are close to simple, such as quasisimple and almost-simple groups. In this article, we consider the ‘natural module’ for groups of Lie type in defining characteristic, simple modules for various low-rank Lie type groups in small, non-defining characteristics, alternating and symmetric groups, and nine of the sporadic groups.

On natural modules, we have the following result.

Theorem 1.1 *Let q be a power of a prime, and let G be a finite group of Lie type. If G is a classical group, suppose that it is one of the groups $\mathrm{SL}_n(q)$ or $\mathrm{Sp}_{2n}(q)$ for $n \geq 3$, $\mathrm{SU}_n(q)$ for $n \geq 6$, $\Omega_{2n}^+(q)$ or $\Omega_{2n}^-(q)$ for $n \geq 4$, or $\Omega_{2n+1}(q)$ for $n \geq 3$. If G is of exceptional type, suppose that it is one of the groups ${}^3D_4(q)$, ${}^2G_2(q)$, ${}^2F_4(q)$, $G_2(q)$, $F_4(q)$, $E_6(q)$, ${}^2E_6(q)$, $E_7(q)$, and $E_8(q)$. Then the natural module for G is non-algebraic.*

For the classical groups, the natural module is clear, and we will define what we mean by natural module for the exceptional groups in Section 5, but it is generally the non-trivial simple module of smallest dimension.

For the Lie-type groups not mentioned in this theorem ($\mathrm{SL}_2(q)$, $\mathrm{Sp}_4(q)$, $\mathrm{SU}_n(q)$ for $n \leq 5$ and $\mathrm{Suz}(q)$), it is only known for $\mathrm{SL}_2(q)$ that the natural module is actually algebraic: this is a result of Alperin [2] for q even, and unpublished work of Kovács [27] for q odd. Since the q odd case remains unpublished, here we will provide a short proof using the theory of tilting modules, from which it is possible to calculate any tensor product of simple modules for $\mathrm{SL}_2(q)$.

Theorem 1.2 (Alperin, Kovács) *Let p be a prime, let $n \geq 1$ be an integer, and let k be a field of characteristic p . If $G = \mathrm{SL}_2(p^n)$, then all simple kG -modules are algebraic.*

This theorem has the following obvious corollary, using the fact that the natural module for $\mathrm{SL}_2(q)$, hence for $\mathrm{GL}_2(q)$, is algebraic, and restrictions of algebraic modules are algebraic.

Corollary 1.3 *Let p be a prime, and let k be a finite field. If G is a finite group, and M is a 2-dimensional kG -module, then M is algebraic.*

Turning to non-defining characteristic, our results focus on the primes 2 and 3. If $\mathrm{char} k = 2$, then we analyze the groups $\mathrm{PSL}_2(q)$, $\mathrm{PSL}_3(q)$ (for $q \equiv 3 \pmod{4}$) and $\mathrm{PSU}_3(q)$ (for $q \equiv 1 \pmod{4}$). In other words, we analyze the cases where the Sylow 2-subgroups are either dihedral or semidihedral. (In the next two theorems, we take an algebraically closed field, purely so that we are guaranteed the existence of all simple modules that ‘should’ exist.)

Theorem 1.4 *Let k be an algebraically closed field of characteristic 2.*

- (i) *If $G = \mathrm{PSL}_2(q)$, then all simple kG -modules are algebraic if and only if $q \not\equiv 7 \pmod{8}$. If $q \equiv 7 \pmod{8}$, then the two $(q-1)/2$ -dimensional simple modules are non-algebraic.*

- (ii) If $G = \text{PSL}_3(q)$, then all simple kG -modules are algebraic if $q \equiv 3 \pmod 8$. If $q \equiv 7 \pmod 8$, then the two non-trivial simple modules in the principal 2-block are non-algebraic.
- (iii) If $G = \text{PSU}_3(q)$, then all simple kG -modules are algebraic if $q \equiv 1 \pmod 4$.

Theorem 1.5 *Let k be an algebraically closed field of characteristic 3, and let G be a finite simple group such that $C_3 \times C_3$ is a Sylow 3-subgroup of G .*

- (i) *If G is a finite group of Lie type, an alternating group, or one of the sporadic groups M_{22} or HS , then all simple kG -modules in the principal 3-block are algebraic.*
- (ii) *If G is one of the groups M_{11} or M_{23} , then there are non-algebraic simple modules in the principal 3-block.*

Notice that, if G is a finite group of Lie type with elementary abelian Sylow 3-subgroups, and k is a field of characteristic 3, then all simple kG -modules in the principal block are algebraic. In characteristic 5, however, this does not occur, thanks to the following proposition.

Proposition 1.6 *Let k be a field of characteristic $p > 2$, let G be a finite group with non-cyclic Sylow p -subgroups, and let M be an indecomposable kG -module with $p \nmid \dim M$. If M lies on the second row of its component of the Auslander–Reiten quiver, then M is non-algebraic.*

For $F_4(2)$ in characteristic 5, the simple module M of dimension 875823 lies on the second row of its component of the Auslander–Reiten quiver [22], and the Sylow 5-subgroups of $F_4(2)$ are non-cyclic, so M is non-algebraic.

One can use Theorem 1.5, together with a standard reduction, to determine exactly which finite groups with Sylow 3-subgroups $C_3 \times C_3$ have algebraic simple modules in the principal 3-block.

Corollary 1.7 *Let G be a finite group with Sylow 3-subgroups of order at most 9, and let k be a field of characteristic 3. Write B for the principal 3-block of kG . All simple B -modules are algebraic if and only if neither M_{11} nor M_{23} is a composition factor of G .*

For symmetric and alternating groups, we study simple modules in blocks with defect group $C_p \times C_p$, and prove the following result.

Theorem 1.8 *Let p be one of 2, 3 and 5, and let k be a field of characteristic p . Let G be either an alternating group \mathfrak{A}_n or a symmetric group \mathfrak{S}_n for some n . If B is a block of kG with defect group $C_p \times C_p$, then all simple B -modules are algebraic.*

The reason that this result cannot be extended further is that it is difficult to prove whether the simple modules in the principal p -block of $k\mathfrak{S}_{2p}$ are algebraic for $p \geq 7$, but it indeed appears possible that this is true.

Finally, in Section 8 we consider nine of the sporadic groups: the five Mathieu groups, HS , J_2 , Suz and He . In some cases, for example, HS in characteristic 3, all

simple modules are algebraic, and we refer to that section for the specific results that we achieve in this direction.

In some sense, the property of being algebraic seems to be often held by simple modules in ‘small’ blocks, in the sense that the defect group is abelian, or dihedral for example. The question of a module being algebraic seems reasonable to ask in the light of the connections with the homological structure of a block [9], and studying this and related properties is likely the beginning of a deep theory connecting the tensor and homological structure of blocks of finite groups.

The organization of this paper is as follows: the next section includes the quoted and preliminary results that we need, and Section 3 is concerned with $SL_2(q)$. The next two sections deal with classical and exceptional groups, and then Lie-type groups in non-defining characteristic in Section 6. The alternating and symmetric groups are considered in Section 7, and the final section, Section 8, is on the sporadic groups.

Throughout this article, all groups are finite and all modules are finite dimensional, unless otherwise specified.

2 Quoted Results and Preliminaries

In this section we summarize results gathered from the literature, together with a few other results needed in the sections to come.

The basic properties of algebraic modules may be found in [18] for example: being algebraic is closed under direct sums, tensor products, taking summands, induction, restriction, Green correspondence and taking sources. This makes the first result trivial.

Proposition 2.1 *Let M be a kG -module, and let H be a p' -index subgroup of G . We have that M is algebraic if and only if $M \downarrow_H$ is algebraic.*

The next three results are in the literature, and we quote them.

Theorem 2.2 [8] *Let G be a group with abelian Sylow 2-subgroups, and let k be a field of characteristic 2. All simple kG -modules are algebraic.*

Theorem 2.3 [9, Theorem B] *Let G be a finite group and let k be a field of characteristic p . If M is a non-periodic, algebraic kG -module, then $\Omega^i(M)$ is non-algebraic for all $i \neq 0$. If M is a periodic, algebraic kG -module then $\Omega^i(M)$ is algebraic for all $i \in \mathbb{Z}$.*

Theorem 2.4 *Let G be a finite group, and let k be a field of characteristic p . If B is a block of kG with cyclic or Klein-four defect group, then all simple B -modules are algebraic.*

Proof For blocks with cyclic defect group this follows from the simple fact that there are only finitely many indecomposable modules of a group with cyclic vertex, and for Klein-four defect group this result is in [11]. \square

The next easy observation is very useful when dealing with sporadic groups in particular, as it allows us to prove that many modules are non-algebraic almost immediately.

Proposition 2.5 (V_4 restriction test) *Let G be a finite group, let Q be a Klein-four subgroup of G , and let k be an algebraically closed field of characteristic 2. If M is a kG -module such that $M \downarrow_Q$ contains non-trivial odd-dimensional indecomposable summands, then M is non-algebraic.*

Proof By [6], the odd-dimensional indecomposable kG -modules are $\Omega^i(k)$ for $i \in \mathbb{Z}$, and $i \neq 0$ for them to be non-trivial. By Theorem 2.3 these modules are non-algebraic, and hence M is non-algebraic. \square

As a remark, by [3, Theorem 3.4] this result holds with Q replaced by any dihedral 2-group.

Proposition 2.6 [9] *Let $k = \mathbb{F}_3$ and let $G = C_3 \times C_3$. If M is an absolutely indecomposable kG -module of dimension either 3 or 6, then M is periodic if and only if it is algebraic.*

We remind the reader of [9, Conjecture E], which motivates some of the expected results in this article.

Conjecture 2.7 *Let k be a finite field of characteristic p , and let $G = C_p \times C_p$, where p is an odd prime. If M is an absolutely indecomposable kG -module of dimension a multiple of p , then M is algebraic if and only if it is periodic.*

Let $T(M)$ denote the (infinite-dimensional) module

$$T(M) = \bigoplus_{n \geq 1} M^{\otimes n},$$

and let $\mathcal{T}(M)$ denote the set of isomorphism classes of indecomposable summands of $T(M)$.

Lemma 2.8 *Let G be a finite group, and let k be a field of characteristic p . Let M be an indecomposable kG -module. If $M^{\otimes 2} = M \oplus X$, where X is algebraic, then M is algebraic.*

Proof Let $\mathcal{X} = \mathcal{T}(X)$, and let \mathcal{M} denote the (finite) set of isomorphism classes of indecomposable summands of $M \otimes A$, as A runs through \mathcal{X} , together with M itself. We claim that $\mathcal{M} = \mathcal{T}(M)$. Certainly, as X is a summand of $M^{\otimes 2}$, all modules in \mathcal{M} appear in $\mathcal{T}(M)$. To see the converse, we claim that if $A \in \mathcal{M}$ then all summands of $A \otimes M$ lie in \mathcal{M} . Since $M \in \mathcal{M}$, this will complete the proof. If $A = M$ then $A \otimes M = M \oplus X$, and the claim is true; if $A \in \mathcal{X}$ then again clearly all summands of $A \otimes M$ lie in \mathcal{M} , by construction. If A is a summand of $B \otimes M$, for $B \in \mathcal{X}$, then

$$A \otimes M \mid B \otimes M \otimes M = B \otimes (M \oplus X) = B \otimes M \oplus B \otimes X.$$

The summands of $B \otimes M$ lie in \mathcal{M} , and the summands of $B \otimes X$ lie in $\mathcal{X} \subseteq \mathcal{M}$, so the proof is complete. \square

Proposition 2.9 *Let p be an odd prime, let k be a field of characteristic p , and let G be a finite group with non-cyclic Sylow p -subgroups. The heart of $\mathcal{P}(k)$, $\text{rad}(\mathcal{P}(k))/\text{soc}(\mathcal{P}(k))$, is non-algebraic.*

Proof Since G has non-cyclic Sylow p -subgroups, k is non-periodic. Since p is odd, k lies on a component Γ of the Auslander–Reiten quiver of type A_∞ . Write $M_1 = k$, and M_{2i+1} for $3 \leq 2i + 1 \leq p$ for the indecomposable modules lying directly above M_1 on Γ (with M_3 the closest to M_1). The modules $\Omega^i(k)$ have dimension $\pm 1 \pmod p$, and it is easy to see that the modules $\Omega^i(M_j)$ have dimension $\pm j \pmod p$; in particular, M_p has dimension a multiple of p . Write Γ' for the component of the Auslander–Reiten quiver containing $\Omega(k)$: write M_2 for the middle term of the almost-split sequence starting in $\Omega(k)$, and M_{2i} for $4 \leq 2i \leq p - 1$ for the modules lying directly above M_2 on Γ' .

Let \mathcal{A}_i denote the almost-split sequence starting in $\Omega(M_i)$, which has middle term $M_{i-1} \oplus M_{i+1}$. By [4, Theorem 3.6], taking the tensor product of \mathcal{A}_1 by M_i yields (up to projectives) \mathcal{A}_i , and so taking middle terms we get

$$M_{i-1} \oplus M_{i+1} = \Omega^0(M_i \otimes M_2),$$

for $1 < i < p$. Finally, since M_p has dimension a multiple of p , $\mathcal{A}_1 \otimes M_p$ is split by [4, Theorem 3.6] again, and so (modulo projectives) $M_p \otimes M_2 = \Omega(M_p) \oplus \Omega^{-1}(M_p)$.

The tensor powers of M_2 contain all M_i for $2 \leq i \leq p$, and hence contain the summands of $M_2 \otimes M_p$. As not both of M_p and $\Omega(M_p)$ can be algebraic, Theorem 2.3 implies that M_2 is non-algebraic, as claimed. \square

Proposition 2.10 *Let p be an odd prime, let k be a field of characteristic p , and let G be a finite group with non-cyclic Sylow p -subgroups. Write E for the heart of $\mathcal{P}(k)$.*

- (i) *If M is a non-algebraic module then $E \otimes M$ is non-algebraic.*
- (ii) *If M is an algebraic module of p' -dimension then $E \otimes M$ contains a non-algebraic summand of dimension prime to p .*

In particular, if $p \nmid \dim M$ then $M \otimes E$ is non-algebraic.

Proof If M is non-algebraic then, since E is self-dual and of dimension prime to p , $M^{\otimes 2}$, which is non-algebraic, is a summand of $(M \otimes E)^{\otimes 2}$. Hence $(M \otimes E)^{\otimes 2}$, so $M \otimes E$, is non-algebraic, proving (i).

For (ii), notice that, since M is algebraic, if all p' -dimensional summands of $E \otimes M$ were algebraic, then the same would be true for $E \otimes M \otimes M^*$ as, if X and Y are absolutely indecomposable modules with one of $\dim X$ and $\dim Y$ a multiple of p , the same is true for all summands of $X \otimes Y$ by [5, Proposition 2.2]. However, as $k \mid M \otimes M^*$, we see that $E \mid E \otimes M \otimes M^*$, a contradiction as E is non-algebraic. This proves (ii). \square

Using this proposition we may easily prove Proposition 1.6 from the introduction. Let M be a kG -module that lies on the second row of its Auslander–Reiten quiver, so that the almost-split sequence with middle term M is

$$0 \rightarrow \Omega^{-1}(N) \rightarrow M \oplus X \rightarrow \Omega(N) \rightarrow 0$$

where X is zero or a projective module. If $p \nmid \dim M$ then, by [4, Theorem 3.6], if E denotes the heart of $\mathcal{P}(k)$, we have that $N \otimes E = M \oplus X$; by Proposition 2.10, this module, hence M , is non-algebraic, as claimed.

We end this section with a proposition, noting that certain types of Morita equivalence preserve algebraicity. (The proof uses the source algebra of a block: see [31] for more information.)

Proposition 2.11 *Let k be a field of characteristic p , and let B and B' be blocks of finite groups, with isomorphic defect groups. If B and B' are splendidly Morita equivalent, then under the induced bijection of simple modules, algebraic modules are sent to algebraic modules, and non-algebraic modules are sent to non-algebraic modules.*

Proof By [30, Remark 7.5], a splendid Morita equivalence induces an isomorphism of source algebras. By [31, Proposition 38.3] (see also the discussion immediately preceding the result in [31]) the sources of simple modules are an invariant of the source algebra, and so the sources of the simple B -modules are isomorphic to the sources of the simple B' -modules, via some isomorphism of defect groups. Since algebraicity is detected in the source of a module, we have the result. \square

3 $SL_2(\mathfrak{q})$

In this section, p is an odd prime, $q = p^n$, $k = \mathbb{F}_q$, and $G = SL_2(q)$. Let $\mathbf{G} = SL_2(\bar{k})$. We embed G inside \mathbf{G} in the standard way, as the fixed points under the map F^n , where $F : \mathbf{G} \rightarrow \mathbf{G}$ is the Frobenius map raising each matrix entry to the p th power.

It is well known that the simple $k\mathbf{G}$ -modules $L(\lambda)$ are labelled by non-negative integers λ , and the simple kG -modules are the restrictions of $L(\lambda)$ for $0 \leq \lambda \leq q$. Let $V_i = L(\lambda)|_G$ for $0 \leq i \leq p - 1$, the so-called *fundamental modules*. By Steinberg’s tensor product theorem, for $\lambda \geq 0$, writing $\lambda = \sum_{i=0}^d j_i p^i$ (for some d), we have that $L(\lambda)$ is the tensor product

$$L(\lambda) \cong \bigotimes_{i=0}^d L(\lambda_i)^{\sigma^i}$$

for $0 \leq \lambda_i \leq p - 1$, where σ is the map on the module category obtained by applying the Frobenius morphism (so that it sends $L(\lambda)$ to $L(p\lambda)$, for example). This tensor product theorem obviously restricts to G , with d replaced by $n - 1$.

We also require tilting modules for $SL_2(\bar{k})$. We will not repeat the definition here, but instead give the properties of them that we need (see [12]). The indecomposable tilting modules, $T(\lambda)$, are parameterized by non-negative integers, with $T(\lambda)$ having

composition factors $L(\lambda)$ and $L(\mu)$ for $\mu < \lambda$, and a tilting module is a sum of the $T(\lambda)$ for various λ . The tensor product of two tilting modules is a tilting module.

We determine some specific tilting modules now, using [16, Lemma 5]. For $0 \leq \lambda \leq p - 1$ we have $T(\lambda) = L(\lambda)$; for $p \leq \lambda \leq 2p - 2$, writing $\lambda = p + \mu$, $T(\lambda)$ is uniserial of length 3, with radical layers $L(p - 2 - \mu)$, $L(\lambda) = L(\mu) \otimes L(1)^\sigma$ and $L(p - 2 - \mu)$; the module $T(2p - 1)$ is simple, and is $L(2p - 1) = L(p - 1) \otimes L(1)^\sigma$. Finally, notice that, since $T(\lambda)$ has composition factors $L(\lambda)$ and $L(\mu)$ for $\mu < \lambda$, a tilting module is determined up to isomorphism by its composition factors.

We now identify the modules V_i : V_0 is the trivial module, V_1 is the 2-dimensional natural module, and $V_i = S^i(V_1)$ is the i th symmetric power of V_1 for $2 \leq i \leq p - 1$. As $S^i(M)$ is a summand of $M^{\otimes i}$ for $i \leq p$, we see that if V_1 is algebraic then so are all fundamental modules. Furthermore, since M is algebraic if and only if the Frobenius twist M^σ is algebraic, we see that if V_1 is algebraic then all simple kG -modules are algebraic, by Steinberg’s tensor product theorem.

The tensor products of $L(\lambda)$ and $L(\mu)$ for $0 \leq \lambda \leq \mu \leq p - 1$ are easy to describe.

Lemma 3.1 *Let $0 \leq \mu \leq \lambda \leq p - 1$.*

(i) *If $\lambda + \mu \leq p - 1$, then*

$$L(\lambda) \otimes L(\mu) = L(\lambda - \mu) \oplus L(\lambda - \mu + 2) \oplus \cdots \oplus L(\lambda + \mu).$$

(ii) *If $\lambda + \mu > p - 1$ and $\lambda < p - 1$ then*

$$\begin{aligned} L(\lambda) \otimes L(\mu) &= L(\lambda - \mu) \oplus L(\lambda - \mu + 2) \oplus \cdots \oplus L(a) \\ &\oplus \begin{cases} L(p - 1) \oplus T(p + 1) \oplus T(p + 3) \oplus \cdots \oplus T(\lambda + \mu) & \mu \text{ even} \\ T(p) \oplus T(p + 2) \oplus \cdots \oplus T(\lambda + \mu) & \mu \text{ odd} \end{cases}, \end{aligned}$$

where $a = 2p - (\lambda + \mu + 4)$.

(iii) *If $\lambda = p - 1$ then*

$$\begin{aligned} L(\mu) \otimes L(p - 1) &= \begin{cases} L(p - 1) \oplus T(p + 1) \oplus T(p + 3) \oplus \cdots \oplus T(p + \mu - 1) & \mu \text{ even} \\ T(p) \oplus T(p + 2) \oplus \cdots \oplus T(p + \mu - 1) & \mu \text{ odd} \end{cases}. \end{aligned}$$

Proof Notice that the decompositions suggested in the lemma are into tilting modules, so it suffices to prove that the composition factors of the tensor product $L(\lambda) \otimes L(\mu)$ match those of the decomposition. For (i) and (ii), this is [16, Lemma 4].

Now suppose that $\mu = 1$ and $\lambda = p - 1$. By [13, Theorem 3.2], $L(1) \otimes L(p - 1) = T(p)$, agreeing with (iii). From this one can easily determine the composition factors of $L(\mu) \otimes L(p - 1)$, and this gives (iii). □

Lemma 3.2 *We have*

$$\begin{aligned} T(p) \otimes L(p - 1) &\cong 2 \cdot (T(p) \oplus T(p + 2) \oplus \cdots \oplus T(2p - 3)) \oplus L(2p - 1) \\ &\cong 2 \cdot L(p - 1) \otimes L(p - 2) \oplus L(p - 1) \otimes L(1)^\sigma. \end{aligned}$$

Consequently, for $0 \leq \lambda, \mu \leq p - 1$, we may express $L(1) \otimes L(\lambda) \otimes L(\mu)$ as a sum $A \oplus B \otimes L(1)^\sigma$, where A and B are sums of tensor products $L(i) \otimes L(j)$ for $i, j \leq p - 1$, and $B = 0$ unless $\lambda = \mu = p - 1$, in which case $B = L(p - 1)$.

Proof Firstly, notice that by Lemma 3.1 and the fact that $L(2p - 1) = L(p - 1) \otimes L(1)^\sigma$, the second and third expressions are equal. Next, since $L(2p - 1) = T(2p - 1)$, all expressions are of tilting modules, so it suffices to check that the composition factors coincide. However, the composition factors of $T(p)$ are $L(1)^\sigma$ and $L(p - 2)$ twice, so the composition factors of $T(p) \otimes L(p - 1)$ are those of $L(p - 1) \otimes L(1)^\sigma$ and $L(p - 1) \otimes L(p - 2)$ twice, which are clearly the same composition factors as the third expression in the lemma.

To see the consequence, note that if $0 < \lambda < p - 1$ then $L(1) \otimes L(\lambda) \otimes L(\mu) = L(\lambda - 1) \otimes L(\mu) \oplus L(\lambda + 1) \otimes L(\mu)$, and similarly for μ , so that $\lambda = \mu = p - 1$, and we have the displayed equation in the lemma. \square

Theorem 3.3 *The module V_1 is algebraic.*

Proof In the case where $d = 1$, the Sylow p -subgroups of G are cyclic, so the result holds: hence we assume that $d > 1$.

We claim that every module in $\mathcal{T}(V_1)$ appears as an indecomposable summand of $M \otimes N$ for simple kG -modules M and N ; let \mathcal{M} denote the set of such summands, and notice that $X \in \mathcal{M}$ if and only if $X^\sigma \in \mathcal{M}$. It suffices to show that if M and N are simple kG -modules then every summand of $V_1 \otimes M \otimes N$ appears in \mathcal{M} . We proceed by induction on $\dim(M \otimes N)$, noting that if this dimension is 1 then the result is trivial.

By Steinberg’s tensor product theorem, we may write $M \otimes N$ as

$$Y = \bigotimes_{i=0}^{n-1} (V_{i,1} \otimes V_{i,2})^{\sigma^i}.$$

Write $Y = V_{1,1} \otimes V_{1,2} \otimes Y'$, and consider $V_1 \otimes Y$: if $V_{1,1}$ and $V_{1,2}$ are not both V_{p-1} then $V_1 \otimes V_{1,2} \otimes V_{1,2}$ is a sum of tensor products of two fundamental modules, by Lemma 3.2, and hence $V_1 \otimes Y$ is a sum of tensor products of two simple modules, as required. Hence both $V_{1,1}$ and $V_{1,2}$ are V_{p-1} , and $V_1 \otimes V_{1,1} \otimes V_{1,2}$ can be written as the sum $A \oplus B \otimes V_1^\sigma$, where A and B are non-zero sums of tensor products of fundamental simple modules. Hence $V_1 \otimes Y = A \otimes Y' \oplus V_1^\sigma \otimes B \otimes Y'$; the module $A \otimes Y'$ is a sum of tensor products of two simple modules, so all summands of it lie in \mathcal{M} , and $(V_1^\sigma \otimes B \otimes Y')^{\sigma^{-1}}$ is a tensor product $V_1 \otimes M' \otimes N'$ (as $B = V_{p-1}$) with $\dim(M' \otimes N') < \dim(M \otimes N)$, whence all summands of it lie in \mathcal{M} by induction. As $X \in \mathcal{M}$ if and only if $X^\sigma \in \mathcal{M}$, all summands of $V_1^\sigma \otimes B \otimes Y'$ lie in \mathcal{M} , as needed. \square

4 Natural Modules for the Classical Groups

We begin by recalling Jennings’s theorem on the group algebras of p -groups. Let P be a finite p -group. Define the *dimension subgroups*

$$\Delta_i = \Delta_i(P) = [P, \Delta_{i-1}(P)]\Delta_{\lceil i/p \rceil}(P)^p.$$

This is the fastest-decreasing central series whose quotients $\Delta_i(P)/\Delta_{i+1}(P)$ are elementary abelian p -groups.

Theorem 4.1 [21] *Let P be a p -group and $k = \mathbb{F}_p$. Denote by kP the group algebra of P over k .*

(i) *Let $A_i(P)$ be defined by*

$$A_i(P) = \{g \in P : g - 1 \in \text{rad}^i(kP)\}.$$

Then $A_i(P) = \Delta_i(P)$.

(ii) *Suppose that we choose $x_{i,j} \in P$ such that $x_{i,j}\Delta_{i+1}$ form a basis of Δ_i/Δ_{i+1} . Write $X_{i,j} = x_{i,j} - 1$. Then*

$$\prod_{i,j} X_{i,j}^{\alpha_{i,j}}, \quad 0 \leq \alpha_{i,j} \leq p - 1$$

generate kP . Furthermore, if the weight of such a product is defined to be $\sum_{i,j} i\alpha_{i,j}$, then all products of weight i form a basis of $\text{rad}^{i-1}(kP)/\text{rad}^i(kP)$, and all products of weight at most i form a basis for $kP/\text{rad}^i(kP)$.

Now let $P = C_p \times C_p$ be the elementary abelian group of order p^2 , and let $k = \mathbb{F}_p$. Write $M_i = kP/\text{rad}^i(kP)$. Then Jennings' theorem immediately implies the following result.

Proposition 4.2

- (i) *The module kP has $2p - 1$ radical layers.*
- (ii) *The module M_i has dimension $i(i + 1)/2$ if $i \leq p$.*
- (iii) *The module M_i is spanned by all monomials in X and Y of degree at most $i - 1$, for $i \leq p$.*

In particular, (iii) of this proposition implies the next lemma.

Lemma 4.3 *Let $1 \leq i \leq p - 1$ be an integer. Then $S^i(M_2) = M_{i+1}$.*

Proof This is obvious if one remembers that $S^i(M_2)$ is spanned by all monomials of degree i in the basis elements of M_2 , which are $1, X$ and Y . Thus $S^i(M_2)$ is spanned by all monomials in X and Y of degree at most i . □

Finally, recall that if $i < p$ and k is a field of characteristic p , then for any kG -module M the module $S^i(M)$ is a summand of $M^{\otimes i}$.

Proposition 4.4 *The 3-dimensional kP -module M_2 is non-algebraic.*

Proof Firstly, notice that both $S_1 = S^{p-2}(M_2)$ and $S_2 = S^{p-1}(M_2)$ are summands of $M_2^{\otimes(p-2)}$ and $M_2^{\otimes(p-1)}$ respectively, and hence if at least one of S_1 and S_2 is non-algebraic, then M_2 is non-algebraic and the proposition follows.

To see that not both of S_1 and S_2 are algebraic, we simply note that

$$S_2 = \Omega(S_1)^*.$$

Thus by Theorem 2.3, either S_1 or S_2 , and consequently M_2 , is non-algebraic. □

Using this, we easily deal with the special linear groups.

Proposition 4.5 *Let q be a power of a prime p , and let G be the group $SL_n(q)$, where $n \geq 3$. The natural kG -module is non-algebraic.*

Proof The natural module for $SL_n(q)$ restricts to the subgroup $SL_n(p)$ as the natural module for this group, so it suffices to prove the result for this group. Since the natural module for $SL_n(p)$ is algebraic if and only if the natural module for $GL_n(p)$ is algebraic (by Proposition 2.1), it suffices to find a non-algebraic module of dimension n over \mathbb{F}_p for some finite group. This is assured by Proposition 4.4, by taking the sum of a 3-dimensional non-algebraic module with $n - 3$ copies of the trivial module for the group $C_p \times C_p$. □

Before we deal with the other classical groups in general, we need to prove a result about $\mathfrak{A}_8 = \Omega_6^+(2)$.

Lemma 4.6 *Let k be a field of characteristic 2 and let $G = \mathfrak{A}_8 = \Omega_6^+(2)$. The 6-dimensional natural kG -module M (viewing G as $\Omega_6^+(2)$) is non-algebraic.*

Proof Consider the 8-point natural permutation representation of \mathfrak{A}_8 : this is easily seen to be uniserial, with radical layers k , M and k . Let H denote a (transitive) subgroup of \mathfrak{A}_8 isomorphic with $SL_3(2) = PSL_2(7)$, acting on the eight points of the projective line. The restriction of M to H is the heart of the permutation module of $PSL_2(7) = SL_3(2)$ acting on the projective line, and this is the sum of the two 3-dimensional simple modules. Since these are both non-algebraic by Proposition 4.5, we see that M is non-algebraic, as claimed. □

(As a remark, it can easily be shown that all non-trivial simple modules in the principal 2-block of \mathfrak{A}_8 are non-algebraic.)

Using the well-known theory of alternating forms (see [32, Chapter 3] for example) it is easy to see that for $n \geq 4$ there is a subgroup H of $G = Sp_{2n}(q)$ isomorphic with $Sp_6(q)$, such that the natural module for G restricts to the sum of the natural module for H and a $(2n - 6)$ -dimensional trivial module. For $G = Sp_6(q)$, one may embed $H = SL_3(q)$ as block-diagonal matrices, with $h \in H$ as the top 3×3 -matrix. In this case, the restriction of the natural module for G is the sum of the natural module for H and its dual, so that the natural module for $Sp_{2n}(q)$ is non-algebraic.

The case of orthogonal groups is similar: for $G = \Omega_{2n}^\pm(q)$ with $n \geq 4$ and q even, there is a copy of $H = \Omega_6^+(2)$ embedded so that the natural module for G restricts to the sum of the natural module for H and a $(2n - 6)$ -dimensional trivial module. For odd q , we reduce to $\Omega_6^+(q)$ in the same way as for even q , and here again see a diagonally embedded copy of $SL_3(q)$ in the same way. Hence for orthogonal groups $\Omega_{2n}^\pm(q)$, $n \geq 4$, the natural module is non-algebraic. The case of $\Omega_{2n+1}(q)$, $n \geq 3$, is very similar, and is omitted.

For unitary groups $SU_n(q)$, things are similar, but only when $n \geq 6$. Consider $SU_n(q)$ as those matrices A in $SL_n(q^2)$ such that $(\bar{A})^t = A^{-1}$, where \bar{A} denotes the matrix obtained from A by raising each entry to the q th power. One notes that $SU_6(q) \cap SL_6(q)$ is the set of all matrices satisfying $A^t = A^{-1}$, so that it is orthogonal for odd q and symplectic for even q . As we showed that, for both of these, the natural module is non-algebraic, this completes the statement.

Collating these statements, we have the following.

Proposition 4.7 *Suppose that k is an algebraically closed field of characteristic p , let $q = p^a$, and let $G = Sp_{2n}(q)$, $SU_{n+3}(q)$, $SO_{2n}^+(q)$ or $SO_{2n+2}^-(q)$ (q odd or even), or $SO_{2n+1}(q)$ (q odd), where $n \geq 3$. The natural kG -module is non-algebraic.*

In the case of $Sp_4(q)$ it is not known in general what happens. For $q = 2$, $G = Sp_4(q) \cong \mathfrak{S}_6$, and in this case it is true that all simple kG -modules are algebraic, as we shall see in Theorem 7.3. However, in general it is not clear whether the natural module is algebraic. (Note that the Suzuki groups $Suz(2^{2n+1})$ are likely to follow the same pattern as $Sp_4(2^n)$.)

For unitary groups, we are interested in $SU_n(q)$ for $n = 3, 4, 5$. The group $SU_3(2)$ is soluble, so the natural module is algebraic, and the natural module for $SU_3(3)$ is algebraic, by a computer-based proof. For $q \geq 4$ however, it is not clear whether the natural module for $SU_3(q)$ is algebraic, even using a computer.

For $n = 4$, the natural module for $SU_4(2)$ is non-algebraic, as there is a conjugacy class of V_4 subgroups such that the natural module restricts to the sum of a 1- and 3-dimensional module, and so is non-algebraic by the V_4 restriction test (Proposition 2.5). By embedding $SU_4(2)$ into $SU_n(q)$, where q is even and $n = 4, 5$, we see that the natural module is non-algebraic for even q and all $n \geq 4$. It appears as though this statement also holds for odd q , but as of yet the author does not have a proof.

5 The Natural Module for the Exceptional Groups

In this section we will prove that the natural module for the exceptional groups of Lie type is non-algebraic in all cases apart from the Suzuki groups. The strategy in the larger-rank cases is as follows: for an algebraic group G , find a subgroup H such that the natural module, restricted to H , has a simple summand that is known to be non-algebraic by earlier results. For small-rank groups, we use some direct computation and some general results on restricting modules. We use the notation from [28] for the weights of algebraic groups.

Proposition 5.1 *If G is the group ${}^2G_2(q)$, where q is an odd power of 3, then the 7-dimensional natural module is non-algebraic.*

Proof Let G be the group ${}^2G_2(3) = SL_2(8) \rtimes C_3$, and let M denote the natural kG -module, where k is a splitting field of characteristic 3. This group has a 9-point permutation representation, and the corresponding permutation module X is uniserial, with radical layers k , M and k .

Let P be a Sylow 3-subgroup of G . It is generated by an element x of order 9, which generates a Sylow 3-subgroup of $SL_2(8)$, and y of order 3, acting non-trivially on $\langle x \rangle$. Let $Q = \langle x^3, y \rangle$, which must act transitively on the nine points; thus we have $X \downarrow_Q = kQ$, and so M restricts to the heart of kQ , which is non-algebraic by Proposition 2.9. Hence M is non-algebraic.

Clearly the natural module for ${}^2G_2(q)$ restricts to ${}^2G_2(3)$ as M , and so we have proved our result. \square

Having dealt with the small Ree groups, we turn our attention to the big Ree groups. The smallest is the Tits group, ${}^2F_4(2)'$, which will require a computer to analyze. Although there is a subgroup of the Tits group isomorphic with $PSL_3(3)$, and the natural module restricts to the 26-dimensional simple module for this group, we will prove in Proposition 6.2 that this module is, in fact, algebraic. Using a computer, we confirm that there are ten conjugacy classes of subgroups of $G = {}^2F_4(2)'$ isomorphic with V_4 , three of which have a normalizer of order 256. One of these is important, and we will need it in the next proposition.

Proposition 5.2 *If G is the group ${}^2F_4(q)'$, where q is an odd power of 2, and M is the 26-dimensional natural module for G , then M is not algebraic.*

Proof There is a conjugacy class of subgroups of $G = {}^2F_4(2)'$ isomorphic with V_4 , and with normalizer of order 256, such that the restriction of M to an element P from this class is

$$M \downarrow_P = 4 \cdot \mathcal{P}(k) \oplus \Omega^2(k) \oplus \Omega^{-2}(k).$$

By Proposition 2.5, $M \downarrow_P$, and hence M , is non-algebraic. Since the natural module for ${}^2F_4(q)$ restricts to the natural module for ${}^2F_4(2)'$, we get the result. \square

In fact, there are five simple modules in characteristic 2 for the Tits group ${}^2F_4(2)'$: three in the principal 2-block and two projective simple modules. The two non-trivial, non-projective, simple modules—of dimensions 26 and 246—are both non-algebraic.

The Suzuki groups are the other twisted groups that only exist in certain circumstances. In this case, nothing is known for any of the groups in characteristic 2; even if we can prove that (say) the natural 4-dimensional module for $Sz(8)$ is non-algebraic, then since $Sz(8)$ is not inside $Sz(32)$ we get nothing directly from this.

We now turn to $G_2(q)$, and the case where q is even needs a separate treatment, given that the natural module has dimension 6 in this case.

Proposition 5.3 *Let G be the group $G_2(q)$, and let M be the 7-dimensional natural module for G , unless $p = 2$, in which case M is the 6-dimensional natural module. Then M is non-algebraic.*

Proof Firstly, assume that q is odd. Let H be the subgroup $SL_3(q) \times C_2$, denoted K_+ in [23, Theorem A]. In [23], Kleidman remarks that M restricts to H as the sum of the trivial and a 6-dimensional simple module. It is clear that the restriction of this 6-dimensional module to $H' \cong SL_3(q)$ is the direct sum of the two (dual) 3-dimensional simple modules. By Proposition 4.5, the 3-dimensional simple modules for $SL_3(q)$ are non-algebraic, and so M is non-algebraic.

Now suppose that q is even. The group $G_2(2) = \text{PSU}_3(3) \times C_2$ has a conjugacy class of Klein-four subgroups with 63 elements, and restricting the 6-dimensional natural module to this class is the sum of two 3-dimensional indecomposable summands; hence by the V_4 restriction test, the natural module for $G_2(2)$ is non-algebraic, and the same therefore holds for $G_2(q)$ by restriction to $G_2(2)$. \square

By considering the embedding of $G_2(q)$ inside ${}^3D_4(q)$, we may prove the next result.

Proposition 5.4 *If G is the group ${}^3D_4(q)$, and M is the 8-dimensional natural module for G , then M is non-algebraic.*

Proof Let H be a maximal subgroup of ${}^3D_4(q)$ isomorphic to $G_2(q)$. If q is odd then the module M restricts as the sum of k and the natural module for H (see the construction in [32, Section 4.6]), so is non-algebraic. For even q this restriction is indecomposable, and one proceeds as in Proposition 5.3: the conjugacy class of Klein-four subgroups with 2,457 members fulfils the requirement of the V_4 restriction test, so the natural module is non-algebraic in this case as well. \square

To work with the rest of the groups, we will restrict simple modules for an algebraic group to a subgroup isomorphic with A_2 , so we need some information about such modules. If G is an algebraic group, H is a subgroup of G isomorphic to A_2 , and M is a module for G such that the restriction of M to H has composition factors only $L(00)$, $L(10)$, $L(01)$ and $L(11)$ (of dimensions 1, 3, 3 and 8 (7 if $p = 3$) respectively), then the only non-trivial extension can occur between $L(00)$ and $L(11)$ by the linkage principle, and so if either $L(10)$ or $L(01)$ is a composition factor then it must be a summand. As the restrictions of these modules to a fixed point subgroup $H \cong \text{SL}_3(q)$ are non-algebraic, the module M for the fixed point subgroup $G = G^F$ is also non-algebraic.

The next group is $F_4(q)$, which has a 26-dimensional natural module. This module is simple unless q is a power of 3, in which case this 26-dimensional module splits as the sum of a 25-dimensional module and the trivial module.

Proposition 5.5 *Let G be the group $F_4(q)$, and let M be the module $L(0001)$, which is 26-dimensional if q is not a power of 3, and 25-dimensional if $3 \mid q$. Then M is not algebraic.*

Proof Let $G = F_4$ and let $H \cong A_2$ be a Levi subgroup corresponding to the two short roots of the root system: the restriction of $L(0001)$ to this subgroup is a sum of one $L(11)$, three $L(10)$, three $L(01)$, and one $L(00)$ if $p = 3$. By the previous remarks, this means that each of the $L(10)$ is a summand of $L(0001) \downarrow_H$, and so, taking fixed points under the Frobenius map, there is a subgroup $H \cong \text{SL}_3(q)$ such that the natural module for H is a summand of $M \downarrow_H$; hence M is non-algebraic, as claimed. \square

We will now deal with the groups $E_6(q)$ and ${}^2E_6(q)$; there is a 27-dimensional natural module of this group in all characteristics.

Proposition 5.6 *If G is either $E_6(q)$ or ${}^2E_6(q)$, and M is the 27-dimensional natural module, then M is non-algebraic.*

Proof Let $G = E_6$, and let H be a subgroup of E_6 isomorphic with F_4 . By [28, Proposition 2.5], the restriction of the 27-dimensional natural module for G to H is the sum of the natural module $L(0001)$ and the trivial module, for all $p \neq 3$. In the case where $p = 3$, we use a different subgroup, namely the one isomorphic to C_4 , which only exists for odd primes. By [28, Proposition 2.5] again, the restriction to this C_4 subgroup is $L(0100)$.

Hence we consider the module $L(0100)$ for $G_1 = C_4$. Let $H_1 \cong A_2$ be the Levi subgroup corresponding to the A_2 on the end of the Dynkin diagram. The restriction of $L(0100)$ to this subgroup is a sum of one $L(11)$, three $L(10)$, three $L(01)$, and two $L(00)$ (only one if $p > 3$). Hence $L(10)$ is a summand of $L(0100) \downarrow_{H_1}$.

Taking fixed points under the appropriate Frobenius map, and noting that $F_4(q)$ and $Sp_8(q)$ (q odd) are contained in ${}^2E_6(q)$, we see that for all q the module M is non-algebraic since the restriction of M to $F_4(q)$ or $Sp_8(q)$ is non-algebraic (the latter case because its restriction to the $SL_3(q)$ -Levi is non-algebraic). □

To deal with the groups $E_7(q)$ and $E_8(q)$, we use the fact discussed above for modules for A_2 .

Proposition 5.7 *Let G be either $E_7(q)$ and $E_8(q)$, and let M denote the natural module, of dimension 56 and 248 respectively. Then M is non-algebraic.*

Proof We use the various tables from [28]. If a module N for A_2 has composition factors $L(00)$, $L(10)$, $L(01)$ and $L(11)$, then we say that N has ‘the desired composition factors’. This will complete the proof for $E_7(q)$ and $E_8(q)$ by taking fixed points under the appropriate Frobenius map.

If $G = E_7$, then by [28, Proposition 2.3] there is a subgroup isomorphic with a central product A_2A_5 , and restricting the 56-dimensional simple module to the A_2 factor has the desired composition factors.

Now let $G = E_8$. By [28, Proposition 2.1], there is a subgroup that is a central product A_2E_6 , and restricting the Lie algebra module to the A_2 factor has the desired composition factors. □

One should note that the same argument, using [28, Proposition 2.1], proves that the kG -module corresponding to the Lie algebra module is non-algebraic for G of type F_4 , E_6 and E_7 as well.

6 Non-Defining Characteristic Groups

In this short section we use a mixture of theoretical and computational techniques to prove results in characteristics 2 and 3. We use known theorems about the fact that the sources of simple modules in non-defining characteristic are ‘generic’, in the sense that they do not depend on the underlying field but only on congruences. We start with the following easy proposition.

Proposition 6.1 *Let G be the group $\mathrm{PSL}_2(q)$ for q odd, and let k be a field of characteristic 2. If $q \equiv -1 \pmod{8}$ then the non-trivial simple modules in the principal block of kG are non-algebraic, and in all other cases all simple modules are algebraic.*

Proof We may assume that $\mathbb{F}_4 \subseteq k$, so that k is a splitting field for G . If $q \equiv 3, 5 \pmod{8}$, then the Sylow 2-subgroup of G is Klein-four: hence all simple kG -modules are algebraic by Theorem 2.4. If $q \equiv 1 \pmod{4}$ then by [14, Theorem 1] the source of a non-trivial simple module in $B_0(kG)$ is 2-dimensional, hence algebraic by Corollary 1.3. If $q \equiv 3 \pmod{4}$ then by [14, Theorem 3] the source of a non-trivial simple module in $B_0(kG)$ has dimension $2^{a-1} - 1$, where 2^a is the order of a Sylow 2-subgroup of G . Since $2^{a-1} - 1$ is odd and greater than 1, and dihedral 2-groups have no non-trivial odd-dimensional indecomposable modules that are algebraic by [3, Theorem 3.4], this module cannot be algebraic, as claimed.

It is easy to see that for $\mathrm{PSL}_2(q)$ there is a unique block with non-cyclic defect group, and so the proof is complete. \square

The remaining simple group with dihedral Sylow 2-subgroups is \mathfrak{A}_7 . By [14, Theorem 5], the three simple modules in the principal 2-block have source either trivial or 2-dimensional, so these are algebraic, just as in the previous proposition. As the non-principal 2-block has Klein-four defect group, we see that all simple $k\mathfrak{A}_7$ -modules are algebraic, where k is a field of characteristic 2.

The two odd central extensions of simple groups with dihedral Sylow 2-subgroups are $3 \cdot \mathfrak{A}_6$ and $3 \cdot \mathfrak{A}_7$: in the former case, there is a faithful 2-block with full defect, and it is splendidly Morita equivalent to the principal 2-block of $\mathrm{PSL}_2(7)$, so there are non-algebraic simple modules by Proposition 2.11; in the latter case, the faithful 2-blocks with full defect have 6-dimensional and 15-dimensional simple modules, both of which are algebraic.

Having dealt with simple groups with dihedral Sylow 2-subgroup, we consider simple groups with semidihedral Sylow 2-subgroup, in characteristic 2. Together with M_{11} , these are $\mathrm{PSL}_3(q)$ and $\mathrm{PSU}_3(q)$, for certain congruences modulo 4. In the case where the Sylow 2-subgroup is semidihedral, work of Erdmann [14] has identified the vertices and sources of the simple modules lying in blocks with full defect, allowing us to prove the following proposition.

Proposition 6.2 *Let G be a simple group with semidihedral Sylow 2-subgroup, and let k be a field of characteristic 2.*

- (i) *If $G \cong \mathrm{PSL}_3(q)$ for $q \equiv 3 \pmod{4}$, then all simple kG -modules in blocks of full defect are algebraic if and only if $q \equiv 3 \pmod{8}$, and if $q \equiv 7 \pmod{8}$ then there is exactly one non-algebraic simple module in each block of full defect.*
- (ii) *If $G \cong \mathrm{PSU}_3(q)$ for $q \equiv 1 \pmod{4}$, then all simple modules in blocks of full defect are algebraic.*
- (iii) *If $G \cong M_{11}$ then all simple kG -modules are algebraic.*

Proof

- (i) Let G act on the projective plane in the standard way, with $q^2 + q + 1$ points, and let H denote a point stabilizer. This is of the form $U \rtimes C$, where C is the

centralizer of an involution and U is a group of odd order. The group C is a quotient of $GL_2(q)$ by a central subgroup of odd order: by Proposition 6.1, if $q \equiv 7 \pmod 8$ then there are non-algebraic simple modules in the principal 2-block of $PSL_2(q)$, and the diagonal automorphism amalgamates these two modules, so that the principal 2-block of kC has a unique non-trivial simple module, and this is non-algebraic.

In [15, (3, 4)], it is shown that the sources of simple modules in blocks of full defect are either trivial source, or uniserial, of dimension $2^{n-3} - 1$, where the power of 2 dividing $|G|$ is $2^n \geq 16$. If $q \equiv 3 \pmod 8$ then $n = 4$, so all simple modules have trivial source, thus are algebraic. If $q \equiv 7 \pmod 8$ then two of the simple modules in the principal block have trivial source, and the other, M , has the property that the restriction to C has as a summand the non-trivial simple module in the principal 2-block of kC : this module is non-algebraic, so M is non-algebraic. In [15, (3,4)] it is also stated that each 2-block of kG contains a simple module whose source is the same as M , completing the proof.

- (ii) This follows immediately from [15, (4.10)], where it is proved that all sources of simple modules in blocks of full defect are of dimension at most 2, so are algebraic by Corollary 1.3.
- (iii) This is proved in Proposition 8.1. □

Let G be the group $SL_3(q)$ for $q \equiv 3 \pmod 4$. The centralizer of an involution is the group $GL_2(q)$, and the defect groups of blocks of $GL_2(q)$ are easy to understand: they are Sylow 2-subgroups of direct products of $GL_d(q)$ for $d \leq 2$, so the blocks are either of full defect or have abelian defect groups. Hence all 2-blocks of G are either of full defect, or of Klein-four or cyclic defect group. Since all simple modules in blocks with the latter two defect groups are algebraic, this means that all simple modules for $PSL_3(q)$ belonging to 2-blocks that are not of full defect are algebraic; in particular, all simple modules for $PSL_3(q)$ are algebraic if $q \equiv 3 \pmod 8$.

Similarly, if G is the group $SU_3(q)$ for $q \equiv 1 \pmod 4$ then the centralizer of an involution is $GU_2(q)$, and again the defect groups are either a Sylow 2-subgroup of G , Klein-four, or cyclic. Hence again, all simple modules lying in 2-blocks of $PSU_3(q)$ (for $q \equiv 1 \pmod 4$) that are not of full defect are algebraic.

We now move on to characteristic 3, and consider simple groups with Sylow 3-subgroup $C_3 \times C_3$.

Proposition 6.3 *Let k be a field of characteristic 3. Let G be one of the simple groups $PSL_3(4)$, $PSU_3(5)$, $PSL_4(2)$, $PSL_5(2)$, $PSU_4(4)$, $PSU_5(4)$, $PSP_4(2)$, $PSP_4(4)$, $PSL_2(q)$ for q a power of 3, \mathfrak{A}_7 , M_{22} and HS . All simple modules in $B_0(kG)$ are algebraic.*

Proof If G is one of $PSP_4(4)$, $PSU_4(4)$ and $PSU_5(4)$, then by [29] there is a splendid Morita equivalence between $B_0(kG)$ and $(C_3 \times C_3) \rtimes D_8$, so that all simple $B_0(kG)$ -modules are algebraic by Proposition 2.11. Similarly, if $G = PSL_5(2)$ then there is a splendid Morita equivalence between $B_0(kG)$ and $(C_3 \times C_3) \rtimes D_8$ by [26], so that all simple $B_0(kG)$ -modules are algebraic.

If G is one of $PSP_4(2) = \mathfrak{S}_6$, \mathfrak{A}_7 or $PSL_4(2) = \mathfrak{A}_8$ then all simple $B_0(kG)$ -modules are algebraic by Theorem 7.3 below. Finally, if G is M_{22} or HS , then all simple $B_0(kG)$ -modules are algebraic by Propositions 8.3 and 8.6, respectively. □

Notice that the two sporadic groups M_{11} and M_{23} are not on this list: by Propositions 8.1 and 8.4 they have non-algebraic simple modules.

We are now in a position to state the main theorem about these groups.

Theorem 6.4 *Let G be a finite group, and suppose that a Sylow 3-subgroup P of G has order at most 9. Let k be a field of characteristic 3. All simple $B_0(kG)$ -modules are algebraic if and only if G does not have a composition factor M_{11} or M_{23} .*

Proof Let G be a finite group of this form. By [19, Section 5], there are normal subgroups H and L of G with $L \leq H$, such that G/H and L are 3'-groups and H/L is a direct product of simple groups and an abelian 3-group. As L is a normal 3'-group, the simple kG -modules and the simple $k(G/L)$ -modules in the principal 3-block are the same, so we may assume that $L = 1$. Also, since G/H is a 3'-group, a simple kH -module M is algebraic if and only if $M \uparrow^G$ is algebraic. Since the simple modules in $B_0(kG)$ are summands of $M \uparrow^G$ for M a simple $B_0(kH)$ -module, we may assume that $G = H$, in which case G is a direct product of simple groups with abelian Sylow 3-subgroups and an abelian 3-group. As $|P| = 9$, if G is the direct product of more than one group then the factors have cyclic Sylow 3-subgroup, whence all simple kG -modules are algebraic, and in the other case G is simple. There is a splendid Morita equivalence between $B_0(kG)$ and $B_0(kH)$, where H is one of $\text{PSL}_3(4)$, $\text{PSU}_3(5)$, $\text{PSL}_4(2)$, $\text{PSL}_5(2)$, $\text{PSU}_4(4)$, $\text{PSU}_5(4)$, $\text{PSp}_4(2)$, $\text{PSp}_4(4)$, $\text{PSL}_2(9)$, \mathfrak{A}_7 , M_{11} , M_{22} , M_{23} , and HS (see [24] and the references contained therein). By hypothesis, $G \not\cong M_{11}$ and $G \not\cong M_{23}$, so by Propositions 2.11 and 6.3 the theorem is proved. \square

7 Alternating and Symmetric Groups

In this section we consider blocks of alternating and symmetric groups having defect groups $C_p \times C_p$. If $p = 2$ then this is the Klein-four group, and all simple modules in such blocks are algebraic by Theorem 2.4. Hence we consider odd primes p : in general, knowing the sources of simple modules in these blocks is a difficult problem, but for small primes this can be done.

The primary tool in this section will be the following theorem of the author [10].

Theorem 7.1 *Let B be a p -block of a symmetric group, and suppose that B has defect group $C_p \times C_p$. If M is any simple B -module then the source of M is isomorphic to the source of a simple module from either $B_0(k\mathfrak{S}_{2p})$ or $B_0(k[\mathfrak{S}_p \wr C_2])$.*

The proof of this theorem uses various combinatorial techniques, and is entirely theoretical. Since all simple $k\mathfrak{S}_p$ -modules are algebraic, we see that all simple $k[\mathfrak{S}_p \wr C_2]$ -modules are algebraic. Hence we get the following corollary.

Corollary 7.2 *If all simple $k\mathfrak{S}_{2p}$ -modules are algebraic, then all simple B -modules are algebraic, where B is any p -block of any symmetric group with defect group $C_p \times C_p$.*

Thus we need to determine whether the simple $k\mathfrak{S}_{2p}$ -modules are algebraic. In general this is unsolved, but for $p = 3$ and $p = 5$ we can prove that they are.

Theorem 7.3 *If $p = 3$ or $p = 5$, and B is a p -block of a symmetric or alternating group with defect group $C_p \times C_p$, then all simple B -modules are algebraic.*

Proof It suffices by the previous corollary to prove that result for \mathfrak{S}_6 and $p = 3$, and \mathfrak{S}_{10} and $p = 5$. Let k be a field of characteristic p . If $p = 3$ then $\mathfrak{A}_6 = \text{PSL}_2(9)$, and all simple $k\text{PSL}_2(9)$ -modules are algebraic by Theorem 1.2, so that all simple $k\mathfrak{S}_6$ -modules are algebraic, proving the theorem.

Now let $p = 5$, and consider \mathfrak{A}_{10} . The 8-dimensional simple $k\mathfrak{A}_{10}$ -module has 8-dimensional source A_2 , and there are twenty-six different non-projective, indecomposable modules appearing in tensor powers of A_2 : twenty-two periodic modules, A_2 , the trivial module $k = A_1$, and the sources A_3 and A_7 of the simple modules of dimensions 28 and 56, which are hence also algebraic.

The 34-dimensional simple module has a 9-dimensional source A_4 , and all nine indecomposable summands of tensor powers of A_4 lie in $A_4^{\otimes 2}$. The two 35-dimensional simple, periodic, modules have 10-dimensional sources A_5 and A_6 , and there are ten different non-projective indecomposable summands in their tensor powers. (These modules are swapped by the outer automorphism of \mathfrak{A}_{10} , so it suffices to check one of them.)

The remaining simple modules in the principal block have dimensions 133, 133 and 217, and the sources have dimensions 8, 8 and 42 respectively. The two 133-dimensional modules are swapped by the outer automorphism of \mathfrak{A}_{10} , and so it suffices to check one of them. There are thirty-four different non-projective summands in $T(A_8)$, with thirty of them periodic.

The last module, A_{10} , has dimension 42. The exterior square $\Lambda^2(A_{10})$ contains a summand M of dimension 47. It is easier to prove that M is algebraic than A_{10} is, and indeed M is algebraic, with 32 different indecomposable modules appearing in $T(M)$, of which the largest has dimension 80. Since $A_{10}^{\otimes 2}$ is a sum of indecomposable modules that also appear in $M^{\otimes 2}$, we see that A_{10} is algebraic, completing the proof for this 5-block. As \mathfrak{A}_{10} has only one block with non-cyclic defect, all simple $k\mathfrak{A}_{10}$ -modules are algebraic, and so all simple $k\mathfrak{S}_{10}$ -modules are algebraic, completing the proof. □

8 Sporadic Groups

In this section we will examine nine sporadic simple groups: the five Mathieu groups, HS , J_2 , Suz and He . Having analyzed the smallest Janko group J_1 in [8], this brings the total to ten.

The results here are almost entirely computer driven, and so we do not provide proofs of our statements that may be checked on the computer. In particular, if a simple module fails the V_4 restriction test, we simply state that it does, without going into details, unless it is non-trivial to prove this, even with a computer.

We deal with the Mathieu groups in detail, then only consider certain other sporadic groups, of particular interest, either because they have a block with abelian defect group or because all simple modules are algebraic for some prime p with non-cyclic Sylow p -subgroups. More detailed proofs are available in the author's thesis [7] for these groups in the majority of cases, and we are brief in our justification.

In this section, the simple modules in the principal p -block are labelled $k = S_1, S_2, \dots, S_n$, ordered by increasing dimension. In one case we consider a non-principal block explicitly, and its simple modules are denoted T_1, \dots, T_m , again ordered by dimension. Information on the simple modules is available in [20].

8.1 The Mathieu Groups

We deal with the five Mathieu groups here.

Proposition 8.1 *Let G be the simple group M_{11} . If $p = 2$ then all simple kG -modules are algebraic, and if $p = 3$ then a simple kG -module is algebraic if and only if it is self-dual, so that there are four simple kG -modules in the principal block that are not algebraic.*

Proof Firstly let $p = 2$. The subgroup $M_9 \rtimes C_2$ has index 55, and the permutation representation on this subgroup is semisimple, the sum of the three modules in the principal block. Hence all simple modules in $B_0(kG)$ are trivial source, so algebraic. The simple modules outside of the principal block are projective, so are also algebraic, proving the result for $p = 2$.

If $p = 3$, then the standard 11-point permutation representation is semisimple, with non-trivial submodule the 10-dimensional self-dual simple module S_4 . The sources of the 10-dimensional simple modules S_5 and S_6 that are not self-dual are $\Omega^{\pm 2}(k)$, and so are not algebraic. If M is one of the 5-dimensional simple modules S_2 and S_3 then $\Lambda^2(M)$ is one of S_4 or S_5 , and as $M^{\otimes 2} = \Lambda^2(M) \oplus S^2(M)$, we see that M is not algebraic. The 24-dimensional simple module S_7 has a 6-dimensional source, and is algebraic since it is periodic, via Proposition 2.6. The only simple module not lying in $B_0(kG)$ is projective, and hence algebraic, completing the proof. \square

Proposition 8.2 *Let G be the simple group M_{12} . If $p = 2$ or $p = 3$ then the 15- and 34-dimensional simple modules (in $B_0(kG)$) are non-algebraic, and all other simple modules are algebraic.*

Proof Firstly suppose that $p = 2$. The non-trivial simple modules lying in $B_0(kG)$ fail the V_4 restriction test, and the non-principal block has Klein-four defect group, so all simple modules in it are algebraic by Theorem 2.4. Hence the result is proved.

Now let $p = 3$. The modules S_2 and S_3 have 10-dimensional sources A_2 and A_3 ; both of these are algebraic, with six different indecomposable modules in their tensor powers. As (with some numbering) $S_7 = \Lambda^2(S_2)$ and $S_8 = \Lambda^2(S_3)$, both of these modules are algebraic.

We turn to the 15-dimensional simples S_4 and S_5 , and the 34-dimensional simple S_6 . We first deal with S_4 : there are two classes of subgroups of order 9 with 220 conjugates, and let Q denote one of these. The restriction of S_4 to Q is

$$S_4 \downarrow_Q = 21 \cdot \mathcal{P}(k) \oplus M \oplus \Omega^{-1}(M),$$

where M is a 15-dimensional, non-periodic indecomposable module. (As S_4 and S_5 are dual, we might need to consider S_5 for this decomposition to hold.) As not both

of M and $\Omega^{-1}(M)$ can be algebraic, S_4 (and hence S_5) is not algebraic. The module S_6 has vertex a Sylow 3-subgroup and 7-dimensional source A_6 ; the module $A_6 \downarrow_Q$ is the heart of the projective indecomposable module for Q , and is hence non-algebraic by Proposition 2.9. This completes the proof. \square

Proposition 8.3 *Let G be the simple group M_{22} . If $p = 2$ then the non-trivial simple modules are non-algebraic, and if $p = 3$ then all simple modules are algebraic.*

Proof Using the V_4 restriction test, we get the result for $p = 2$. For $p = 3$, all non-principal blocks have cyclic defect groups, so it suffices to consider the principal 3-block. In this case, there is a splendid Morita equivalence with the principal block of the Mathieu group $M_{10} = \mathfrak{A}_{6.22}$, all of whose simple modules are algebraic by Theorem 7.3 and Proposition 2.1. This completes the proof, via Proposition 2.11. \square

Proposition 8.4 *Let G be the simple group M_{23} . If $p = 2$ then all non-trivial simple modules in the principal block, possibly apart from that of dimension 252, are non-algebraic. If $p = 3$ then a simple module is algebraic if and only if it does not have dimension 104.*

Proof Using the V_4 restriction test, we get the result for $p = 2$. For $p = 3$, all non-principal blocks have cyclic defect groups, so it suffices to consider the principal 3-block. All simple modules in $B_0(kG)$, apart from the two 104-dimensional simple modules S_3 and S_4 , are trivial-source modules, and hence algebraic.

The modules S_3 and $S_4 = S_3^*$ are non-algebraic: to see this, we note that the source of (say) S_3 is a 5-dimensional module M , and $M^{\otimes 2}$ has $\Omega^2(k)$ as a summand. As $\Omega^2(k)$ is non-algebraic, M is non-algebraic, as claimed. \square

The 252-dimensional module has algebraic restriction to all elementary abelian subgroups of M_{23} , so it is not clear whether this module is algebraic or not. However, the 252-dimensional simple module for M_{24} is non-algebraic, and it restricts to this simple module, which suggests that it is probably not algebraic.

Proposition 8.5 *Let G be the simple group M_{24} . If $p = 2$ then all non-trivial simple modules in the principal block, possibly apart from those of dimension 320 or 1,792, are non-algebraic. If $p = 3$ then a simple module is algebraic if and only if it does not have dimension 770 (there are two simple modules with this dimension).*

Proof For $p = 2$, the V_4 restriction test proves that the non-trivial simple modules, with the exception of the two mentioned, are non-algebraic. Now suppose that $p = 3$: all non-principal blocks have cyclic defect group, so it suffices to consider the seven simple modules in the principal block. The 22-dimensional simple S_2 has vertex a Sylow 3-subgroup P of G , and 4-dimensional source, which has a kernel of order 3. It is easy to check by direct calculation that this module is algebraic. Since $\Lambda^2(S_2) = S_3$, the simple module of dimension 231, and $S_2^{\otimes 2} = \Lambda^2(S_2) \oplus S^2(S_2)$, we see that S_3 is also algebraic. The 483-dimensional simple module S_4 has trivial source (it is a summand of a permutation module of dimension 759), and is hence algebraic. Finally for the algebraic modules, the largest simple module S_7 , of dimension 1243, has a 19-dimensional source M (and vertex P). The tensor square of this module is (up to

projectives) the sum of k and two non-isomorphic 18-dimensional modules, induced from subgroups of order 9 in P . Hence S_7 is algebraic if and only if the 6-dimensional sources N_1 and N_2 of the 18-dimensional indecomposable modules are algebraic. The fact that the N_i are periodic, together with Proposition 2.6, completes the proof that S_7 is algebraic.

We turn our attention to the two (dual) 770-dimensional simple modules S_5 and S_6 , which are non-algebraic. If M denotes the 5-dimensional source of S_5 , then the centre of $Z(P)$ is the kernel of P , and viewed as a module over $C_p \times C_p$, $M^{\otimes 2}$ has as a summand either $\Omega^2(k)$ or $\Omega^{-2}(k)$, so neither S_5 nor $S_6 = S_5^*$ is algebraic. This completes the proof. \square

Again, it is not known whether or not the 320- and 1792-dimensional simple modules are algebraic, although the author considers it unlikely.

8.2 HS , J_2 , Suz and He

Here we consider the Higman–Sims, second Janko, Suzuki and Held sporadic groups. In the case of HS , in characteristic 3 then all simple modules are algebraic, and the same is true of J_2 in characteristics 3 and 5. For the larger two sporadics however, things are more complicated: it appears possible that all simple modules for Suz are algebraic in characteristic 5, as at least the smallest six are, and for He almost all of the simple modules are algebraic for both $p = 3$ and $p = 5$.

Proposition 8.6 *Let G be the simple group HS . If $p = 2$ then a non-trivial simple module is algebraic if and only if it lies outside the principal block. If $p = 3$ then all simple modules are algebraic.*

Proof The result for the prime 2 follows easily from the V_4 restriction test for the principal block and, since the only non-principal block has Klein-four defect group, for those not in the principal block by Theorem 2.4.

Now let $p = 3$. There are seven simple modules in each block of kG with non-cyclic defect group, labelled S_1 to S_7 for the principal block and T_1 to T_7 for the non-principal block. The modules $S_1, S_2, S_3, T_3, T_4, T_5, T_6$ and T_7 are all trivial-source modules, hence algebraic.

There is a subgroup $H \cong M_{22}$ of G , of index 100. All simple kH -modules are algebraic by Proposition 8.3, and the restrictions of S_4, T_1 and T_2 to H are semisimple, hence these modules are algebraic by Proposition 2.1. Also, $S_7 \downarrow_H$ is the sum of simple and projective modules, so S_7 is also algebraic.

This leaves us with S_5 and S_6 . Since $S_6 = \Lambda^2(T_1)$, and $T_1^{\otimes 2} = \Lambda^2(T_1) \oplus S^2(T_1)$, this is algebraic. Finally, S_5 has a 10-dimensional source M , and $M^{\otimes 2}$ is the sum of a trivial module, nine free modules, and three 6-dimensional periodic indecomposable modules, which are algebraic by Proposition 2.6. As $M^{\otimes 2}$ is algebraic, M is algebraic so S_5 is algebraic, completing the proof. \square

There are unfortunately no results on the (non-trivial) simple modules in the principal 5-block of HS .

Proposition 8.7 *Let G be the Janko group J_2 . If $p = 2$ then all simple modules outside the principal block are algebraic, and all non-trivial simple modules in the principal block are non-algebraic. If $p = 3$ or $p = 5$ then all simple modules are algebraic.*

Proof If $p = 2$ then the V_4 restriction test is enough to prove the result for the principal block and, since the only non-principal block has Klein-four defect group, for those simple modules not in the principal block by Theorem 2.4.

If $p = 3$, then all simple modules lying outside the principal block lie in blocks with cyclic defect groups, so that they are all algebraic. There are eight simple modules lying in $B_0(kG)$, labelled as usual S_1 to S_8 . Apart from the trivial module $k = S_1$ and the 133-dimensional module S_8 , the other six come in pairs, of dimension 13, 21 and 57, each defined over \mathbb{F}_9 . Because of this, proving (for example) that S_2 is algebraic proves that S_3 is algebraic. We have firstly that

$$S_2 \otimes S_2 = k \oplus S_4 \oplus S_6 \oplus T_2,$$

where T_2 is the 90-dimensional simple module in the second block. If S_2 is algebraic then so are S_4 and S_6 , and by applying the Frobenius morphism on \mathbb{F}_9 , we see that S_3 , S_5 and S_7 are also algebraic. Finally, $S_2 \otimes S_3 = S_8 \oplus T_1$ (where T_1 is the 36-dimensional simple module in the second block), so that S_8 is also algebraic. Thus it remains to prove that S_2 is algebraic. Let A_2 denote the source of S_2 , which is simply $S_2 \downarrow p$. There are nine non-projective modules in $\mathcal{T}(A_2)$, so it is algebraic.

Finally, suppose that $p = 5$. There are six simple modules in the principal block of kG , and all non-principal blocks have cyclic defect groups. The 14-dimensional simple module S_2 has indecomposable restriction to P , and hence its source A_2 is 14-dimensional. It is algebraic, and there are twenty-one non-projective modules in $\mathcal{T}(A_2)$. Using this we may prove that all simple modules are algebraic: we have

$$S_2 \otimes S_2 = k \oplus S_2 \oplus S_3 \oplus T_1 \oplus T_2,$$

where T_1 and T_2 are the two simple modules lying in the second block (with defect 1), of dimensions 70 and 90 respectively. Hence S_3 is algebraic, and in addition

$$S_3 \otimes S_3 = k \oplus S_2 \oplus S_3 \oplus S_4 \oplus S_5 \oplus S_6 \oplus T_2,$$

so that all simple modules in $B_0(kG)$ are algebraic, as required. □

Proposition 8.8 *Let G be the Suzuki sporadic group Suz . If $p = 2$ then all non-trivial simple modules in the principal block are non-algebraic. If $p = 3$ then all non-trivial simple modules in the principal block are non-algebraic, except possibly for the 8436- and 32967-dimensional modules, and all simple modules outside of the principal block are algebraic. If $p = 5$ then the smallest six simple modules in the principal block are algebraic.*

Proof Firstly, let $p = 2$. We apply the V_4 restriction test, but since the simple modules for Suz are quite large, we describe completely the restrictions involved.

Let Q be a representative from the conjugacy class of V_4 subgroups with 1,216,215 members. We have decompositions of:

$$S_2 \downarrow_Q = 8 \cdot k \oplus \Omega^2(k) \oplus \Omega^{-2}(k) \oplus 20 \cdot \mathcal{P}(k) \oplus X_1;$$

$$S_4 \downarrow_Q = 6 \cdot k \oplus 4 \cdot \Omega^2(k) \oplus 4 \cdot \Omega^{-2}(k) \oplus 24 \cdot \mathcal{P}(k);$$

$$S_5 \downarrow_Q = 8 \cdot \Omega(k) \oplus 8 \cdot \Omega^{-1}(k) \oplus 2 \cdot \Omega^3(k) \oplus 2 \cdot \Omega^{-3}(k) \oplus 124 \cdot \mathcal{P}(k);$$

$$S_7 \downarrow_Q = 4 \cdot k \oplus 4 \cdot \Omega(k) \oplus 4 \cdot \Omega^{-1}(k) \oplus \Omega^4(k) \oplus \Omega^{-4}(k) \oplus 148 \cdot \mathcal{P}(k);$$

$$S_8 \downarrow_Q = 8 \cdot k \oplus 16 \cdot \Omega(k) \oplus 16 \cdot \Omega^{-1}(k) \oplus 4 \cdot \Omega^2(k) \oplus 4 \cdot \Omega^{-2}(k) \\ \oplus 4 \cdot \Omega^3(k) \oplus 4 \cdot \Omega^{-3}(k) \oplus 808 \cdot \mathcal{P}(k);$$

$$S_9 \downarrow_Q = 4 \cdot k \oplus 4 \cdot \Omega(k) \oplus 4 \cdot \Omega^{-1}(k) \oplus \Omega^4(k) \oplus \Omega^{-4}(k) \oplus 1116 \cdot \mathcal{P}(k);$$

$$S_{10} \downarrow_Q = 24 \cdot k \oplus 16 \cdot \Omega^2(k) \oplus 16 \cdot \Omega^{-2}(k) \oplus 4 \cdot \Omega^4(k) \oplus 4 \cdot \Omega^{-4}(k) \oplus 1184 \cdot \mathcal{P}(k);$$

$$S_{11} \downarrow_Q = 8 \cdot k \oplus 6 \cdot \Omega(k) \oplus 6 \cdot \Omega^{-1}(k) \oplus 6 \cdot \Omega^3(k) \oplus 6 \cdot \Omega^{-3}(k) \oplus 2264 \cdot \mathcal{P}(k) \oplus X_2;$$

$$S_{13} \downarrow_Q = 12 \cdot k \oplus 2 \cdot \Omega^2(k) \oplus 2 \cdot \Omega^{-2}(k) \oplus 2596 \cdot \mathcal{P}(k) \oplus X_3.$$

Here X_1 , X_2 and X_3 are the sums of six, seventy-two and forty-four 2-dimensional indecomposable modules respectively. Taking Galois conjugates of the decompositions for S_2 , S_{11} and S_{13} give the decompositions for S_3 , S_{12} and S_{14} , and the decomposition for S_5 is also the decomposition for S_6 . Therefore all non-trivial simple modules in $B_0(kG)$ are non-algebraic, as claimed.

Now let $p = 3$. There are eight conjugacy classes of subgroups of G that are elementary abelian of order 9, and let Q be a representative from the class with 38,438,400 members. Write E for the heart of $\mathcal{P}(k)$ and M_1 for the module $\text{soc}^2(\mathcal{P}(k))$. There exist M_2 , a 9-dimensional, self-dual, non-periodic indecomposable module, and M_3 , a self-dual, 37-dimensional indecomposable module, such that the restrictions of the simple modules S_i in the principal block are as follows:

$$S_2 \downarrow_Q = E \oplus \Omega(M_1) \oplus \Omega^{-1}(M_1^*) \oplus 3 \cdot \mathcal{P}(k);$$

$$S_3 \downarrow_Q = M_2 \oplus \Omega^2(M_1) \oplus \Omega^{-2}(M_1^*) \oplus 5 \cdot \mathcal{P}(k);$$

$$S_4 \downarrow_Q = k \oplus M_2 \oplus \Omega^2(M_1) \oplus \Omega^{-1}(M_1^*) \oplus 28 \cdot \mathcal{P}(k);$$

$$S_5 \downarrow_Q = \Omega(M_1) \oplus \Omega^{-1}(M_1^*) \oplus \Omega^3(M_1^*) \oplus \Omega^{-3}(M_1) \oplus 41 \cdot \mathcal{P}(k);$$

$$S_6 \downarrow_Q = E \oplus \Omega(M_1) \oplus \Omega^{-1}(M_1^*) \oplus 68 \cdot \mathcal{P}(k);$$

$$S_7 \downarrow_Q = M_1 \oplus M_1^* \oplus \Omega^3(M_1^*) \oplus \Omega^{-3}(M_1) \oplus \Lambda^2(E) \oplus 209 \cdot \mathcal{P}(k);$$

$$S_8 \downarrow_Q = \Omega(M_1) \oplus \Omega^{-1}(M_1^*) \oplus \Omega^2(M_1) \oplus \Omega^{-2}(M_1^*) \oplus \Omega(M_2) \oplus \Omega^{-1}(M_2) \\ \oplus \Omega^3(M_1) \oplus \Omega^{-3}(M_1^*) \oplus \Omega^4(M_1^*) \oplus \Omega^{-4}(M_1) \oplus 299 \cdot \mathcal{P}(k);$$

$$S_9 \downarrow_Q = M_2 \oplus \Omega^2(M_1) \oplus \Omega^{-1}(M_1^*) \oplus 528 \cdot \mathcal{P}(k);$$

$$S_{11} \downarrow_Q = M_1 \oplus M_1^* \oplus 2 \cdot \Omega(M_1) \oplus 2 \cdot \Omega^{-1}(M_1^*) \oplus \Omega^3(M_1^*) \oplus \Omega^{-3}(M_1) \\ \oplus \Omega(M_2) \oplus \Omega^{-1}(M_2) \oplus 1622 \cdot \mathcal{P}(k);$$

$$S_{12} \downarrow_Q = 2 \cdot M_1 \oplus 2 \cdot M_1^* \oplus \Omega^3(M_1^*) \oplus \Omega^{-3}(M_1) \\ \oplus \Omega^2(k) \oplus \Omega^{-2}(k) \oplus M_3 \oplus 2150 \cdot \mathcal{P}(k);$$

This proves that all non-trivial simple modules S_i are non-algebraic, apart from S_{10} —it appears to be non-algebraic, but the proof of this currently eludes the author—and the 32967-dimensional simple module S_{13} . The latter module has the property that its restriction to any subgroup of order 3 is free, and so the restriction to any subgroup of order 9 is periodic, which makes it difficult to prove that it is non-algebraic. (The size of the module makes a more detailed analysis difficult.)

We turn to the modules outside the principal block. There is only one non-principal block B with non-cyclic defect groups, and in [25], it is proved that B is splendidly Morita equivalent to the principal 3-block of $\text{PSL}_3(4)$; hence all simple B -modules are algebraic by Propositions 2.11 and 6.3.

If $p = 5$, then the 1001-dimensional simple module S_4 is trivial source, hence algebraic. The 143-dimensional simple module S_2 has a 28-dimensional source A_2 , and we have

$$A_2 \otimes A_2 = k \oplus A_2 \oplus X \oplus 29 \cdot \mathcal{P}(k),$$

where X is the (30-dimensional) sum of the permutation modules on the six subgroups of P of order 5. Hence A_2 , and S_2 , are algebraic. The 363-dimensional simple module S_3 has a 13-dimensional source A_3 . There are 37 different non-projective summands of $T(A_3)$, the largest of which has dimension 120; as this number is finite, A_3 is algebraic.

The tensor product of S_2 and S_3 is a module whose only composition factor in the principal block is S_6 , and hence this module is algebraic since it must be a summand of $S_2 \otimes S_3$. Both the 3289-dimensional simple module S_5 and the 11869-dimensional simple module S_6 have 74-dimensional sources A_5 and A_6 , and although they are not isomorphic, they are $\text{Aut}(P)$ -conjugate. Hence S_5 is also algebraic. \square

The reason that simple modules of larger dimension for $p = 5$ were not considered is simply that their dimensions are too high to be easily constructed on a computer. It is possible that the simple modules of larger dimension are also algebraic. As technology improves more simple modules will be easily dealt with; for example, the 41822-dimensional simple module is a composition factor of $\Lambda^2(S_3)$, and this module is *almost* able to be easily constructed on a modern computer without more sophisticated techniques, such as the condensation method.

Proposition 8.9 *Let G be the Held sporadic group He . If $p = 2$ then a simple module is algebraic if and only if it is trivial or lies outside the principal block. If $p = 3$ then a simple module is algebraic if and only if it does not have dimension 6172 or 10879, and if $p = 5$ then the simple modules with dimension 1, 51, 104, 153, 4116, 4249, and 6528 are algebraic.*

Proof First, let $p = 2$. The only non-principal block of kG that is not of defect zero has Klein-four defect, and so all simple modules outside the principal block are algebraic. For the modules in the principal block, the V_4 restriction test proves the result: for all but the largest two simple modules, of dimension 2449, a V_4 subgroup from one of the two conjugacy classes with 437,325 members will work, whereas for the largest two simple modules, a V_4 subgroup from one of the two conjugacy classes with 5,247,900 members is necessary.

We next examine $p = 3$. The 679-dimensional simple module S_2 has a 4-dimensional source A_2 (which has a kernel of order 3): there are seven elements in $\mathcal{T}(A_2)$, so it is easy to prove that it is algebraic. The 3673-dimensional simple module S_4 has 19-dimensional source A_4 , with nine different indecomposable summands in $\mathcal{T}(A_4)$. The 1275- and 6272-dimensional simple modules S_3 and S_6 have trivial source, so are also algebraic.

However, the 6172-dimensional simple module S_5 has a 7-dimensional source A_5 which, when restricted to two of the four maximal subgroups of P (the two that lie in the G -conjugacy class of size 2,332,400) is the heart of the free module. Hence A_5 , and so S_5 , is non-algebraic by Proposition 2.9. Finally, the 10879-dimensional has 16-dimensional source A_7 , and when restricted to the same maximal subgroups as the previous module, is the sum of a free module and the heart of the free module. Hence A_7 , and thus S_7 , is non-algebraic, completing the proof for modules in the principal 3-block.

In [25], it is proved that the non-principal 3-block B of He is splendidly Morita equivalent to its Brauer correspondent, and hence all simple B -modules are trivial source by Proposition 2.11, so algebraic.

Now suppose that $p = 5$. The 51-dimensional simple modules S_2 and S_3 have trivial source, so are algebraic. The 104-dimensional simple module S_4 has a 29-dimensional source A_4 , and there are fifty-eight different indecomposable summands in $\mathcal{T}(A_4)$, the largest of which has dimension 129; thus S_4 is algebraic as well. The exterior square of S_4 has the 4116-dimensional simple module S_{11} as a summand, so this module (which has a 66-dimensional source A_{11}) is also algebraic.

The 153-dimensional simple modules S_5 and S_6 , and the 6528-dimensional simple module S_{13} , all have the same 28-dimensional source A_5 , and $A_5^{\otimes 2}$ is the sum of A_5 , k , the six 5-dimensional permutation modules on subgroups of P of order 5, and a free module: hence A_5 is algebraic by Lemma 2.8, and so S_5 , S_6 and S_{13} are algebraic.

The 4249-dimensional simple module S_{12} has an 8-dimensional source A_{12} , and there are exactly 27 elements of $\mathcal{T}(A_{12})$, the largest of which has dimension 80, so that S_{12} is algebraic. This completes the proof. \square

The remaining simple modules for $p = 5$ are S_7 and S_8 , of dimension 925, S_9 and S_{10} , of dimension 3197, and S_{14} , of dimension 10860. The modules S_7 , S_8 and S_{14} have dimension a multiple of 5, and are non-periodic, so are likely to be non-algebraic, in accordance with Conjecture 2.7: the sources of S_7 and S_8 have dimension 75, and the source of S_{14} has dimension 110, and at the moment we cannot prove whether these are algebraic or not.

The modules S_9 and S_{10} are more complicated: they share a 47-dimensional source A_9 , and we have

$$\begin{aligned} S^2(A_9) &= k \oplus X_1 \oplus A_4 \oplus B_1 \oplus 38 \cdot \mathcal{P}(k) \text{ and } \Lambda^2(A_9) \\ &= X_3 \oplus A_5 \oplus A_9 \oplus A_{11} \oplus 34 \cdot \mathcal{P}(k). \end{aligned}$$

Here, X_i is the sum over all (six) subgroups Q of P of order 5 of the i -dimensional indecomposable module for Q , induced to P , and B_1 is a 118-dimensional simple module. Hence by Lemma 2.8, if B_1 is algebraic then A_9 is algebraic. Taking the

tensor square of B_1 is more difficult, and we just do the symmetric square. This decomposes as

$$S^2(B_1) = k \oplus 2 \cdot X_1 \oplus X_3 \oplus A_4 \oplus B_1 \oplus B_2 \oplus B_3 \oplus B_4 \oplus 256 \cdot \mathcal{P}(k).$$

In this decomposition: B_2 is a 61-dimensional indecomposable module, which appears as a summand in $S^2(A_{11})$, hence algebraic; B_3 is a sum of four 10-dimensional periodic modules, each of which is easily proved to be algebraic; and B_4 is a 222-dimensional indecomposable module. The module B_4 is much more difficult to analyze, and at present we cannot get any information about this module. However, with the plethora of algebraic modules appearing in its tensor powers, the author believes that S_9 and S_{10} are indeed algebraic.

Acknowledgements The author would like to thank the referee for requiring him to redo all of his computer calculations, which led to several typos in the results being corrected, and generally for improving some aspects of the exposition. In addition, he pointed out some errors in the calculations in [8], which have been added to the author's homepage.

References

1. Alperin, J.: On modules for the linear fractional groups. In: Iwahori, N. (ed.) *Finite Groups* (Sapporo and Kyoto, 1974), pp. 157–164. Japan Society for the Promotion of Science, Tokyo (1976)
2. Alperin, J.: Projective modules for $SL(2, 2^n)$. *J. Pure Appl. Algebra* **15**, 219–234 (1979)
3. Archer, L.: On certain quotients of the Green rings of dihedral 2-groups. *J. Pure Appl. Algebra* **212**, 1888–1897 (2008)
4. Auslander, M., Carlson, J.: Almost-split sequences and group rings. *J. Algebra* **103**, 122–140 (1986)
5. Benson, D., Carlson, J.: Nilpotent elements in the Green ring. *J. Algebra* **104**, 329–350 (1986)
6. Conlon, S.: The modular representation algebra of groups with Sylow 2-subgroup $Z_2 \times Z_2$. *J. Aust. Math. Soc.* **6**, 76–88 (1966)
7. Craven, D.: *Algebraic Modules for Finite Groups*. Ph.D. thesis, University of Oxford, Oxford (2008)
8. Craven, D.: Simple modules for groups with abelian Sylow 2-subgroups are algebraic. *J. Algebra* **321**, 1473–1479 (2009) (A list of corrections is available on the author's homepage.)
9. Craven, D.: Algebraic modules and the Auslander–Reiten quiver. *J. Pure Appl. Algebra* **215**, 221–231 (2011)
10. Craven, D.: Sources of simple modules for weight 2 blocks of symmetric groups. *Q. J. Math. Oxford*. doi:[10.1093/qmath/har016](https://doi.org/10.1093/qmath/har016)
11. Craven, D., Eaton, C., Kessar, R., Linckelmann, M.: The structure of blocks with Klein four defect group. *Math. Z.* **268**, 441–476 (2011)
12. Donkin, S.: On tilting modules for algebraic groups. *Math. Z.* **212**, 39–60 (1993)
13. Doty, S., Henke, A.: Decomposition of tensor products of modular irreducibles for SL_2 . *Q. J. Math.* **56**, 189–207 (2005)
14. Erdmann, K.: Principal blocks of groups with dihedral Sylow 2-subgroups. *Commun. Algebra* **5**, 665–694 (1977)
15. Erdmann, K.: On 2-blocks with semidihedral defect groups. *Trans. Am. Math. Soc.* **256**, 267–287 (1979)
16. Erdmann, K., Henke, A.: On Ringel duality for Schur algebras. *Math. Proc. Camb. Philos. Soc.* **132**, 97–116 (2002)
17. Feit, W.: Irreducible modules of p -solvable groups. In: Cooperstein, B., Mason, G. (eds.) *The Santa Cruz Conference on Finite Groups*, pp. 405–412. American Mathematical Society, Providence (1980)
18. Feit, W.: *The Representation Theory of Finite Groups*. North-Holland, Amsterdam (1982)

19. Fong, P., Harris, M.: On perfect isometries and isotypies in finite groups. *Invent. Math.* **114**, 139–191 (1993)
20. Jansen, C., Lux, K., Parker, R., Wilson, R.: *An Atlas of Brauer Characters*. Oxford University Press, New York (1995)
21. Jennings, S.: The structure of the group ring of a p -group over a modular field. *Trans. Am. Math. Soc.* **50**, 175–185 (1941)
22. Kawata, S., Michler, G., Uno, K.: On Auslander-Reiten components and simple modules for finite groups of Lie type. *Osaka J. Math.* **38**, 21–26 (2001)
23. Kleidman, P.: The maximal subgroups of the Chevalley groups $G_2(q)$ with q odd, the Ree groups ${}^2G_2(q)$, and their automorphism groups. *J. Algebra* **117**, 30–71 (1988)
24. Koshitani, S., Kunugi, N.: Broué’s conjecture holds for principal 3-blocks with elementary abelian defect group of order 9. *J. Algebra* **248**, 575–604 (2002)
25. Koshitani, S., Kunugi, N., Waki, K.: Broué’s abelian defect group conjecture holds for the Held group and the sporadic Suzuki group. *J. Algebra* **279**, 638–666 (2004)
26. Koshitani, S., Miyachi, H.: The principal 3-blocks of four- and five-dimensional projective special linear groups in non-defining characteristic. *J. Algebra* **226**, 788–806 (2000)
27. Kovács, L.: Some Indecomposables for SL_2 . Research Report no. 11, Mathematics Research Report Series of the Australian National University (1981)
28. Liebeck, M., Seitz, G.: Reductive subgroups of exceptional algebraic groups. *Mem. Amer. Math. Soc.* **121**(580), vi+111 (1996)
29. Puig, L.: Algèbres de source de certains blocs des groupes de Chevalley. *Astérisque* **181–182**, 221–236 (1990)
30. Puig, L.: *On the Local Structure of Morita and Rickard Equivalences Between Brauer Blocks*. Birkhäuser, Basel (1999)
31. Thévenaz, J.: *G-Algebras and Modular Representation Theory*. Oxford University Press, New York (1995)
32. Wilson, R.: *The Finite Simple Groups*. Springer, London (2009)