



Perverse Equivalences and Broué's Conjecture

David A. Craven

University of Oxford

University of York, 15th March, 2010

Notation and Conventions

Throughout this talk,

- G is a finite group,
- ℓ is a prime,
- K is a field of characteristic ℓ ,
- P is a Sylow ℓ -subgroup of G , and
- Q is a general ℓ -subgroup of G .

I will (try to) use **red** for definitions and **green** for technical bits that can be ignored.

This talk is joint work with Raphaël Rouquier.

From \mathbb{C} -representations to K -representations

Maschke's theorem says that every \mathbb{C} -representation of a finite group G is a sum of simple representations. This is equivalent to $\mathbb{C}G$ being a direct sum of matrix algebras, each of degree that of a representation, with one $\mathbb{C}G$ -module associated to each matrix algebra.

If $K = \bar{\mathbb{F}}_\ell$ (with $\ell \mid |G|$) then this is not true. However, write KG as a sum of indecomposable 2-sided ideals, called **blocks**. Each indecomposable KG -module is associated to a block, but this time **more than one** KG -module is associated to a given block (in general). If an indecomposable module is associated to a block B , then so are all of its composition factors. Hence every block has at least one simple module associated to it.

The number of simple KG -modules in a block B is denoted $\ell(B)$.

Representation Theory is Local

The deepest and most difficult conjectures in representation theory tend to relate the representation theory of G in characteristic ℓ with that of $(\ell\text{-})$ local subgroups $N_G(Q)$, where Q is an ℓ -subgroup of G .

To every block is attached a **defect group** D (an ℓ -subgroup of G up to conjugacy), which ‘controls’ the representation theory of B . The local conjectures are localized further to relate B with a block b of $KN_G(D)$, called the **Brauer correspondent**.

Alperin’s weight conjecture gives a precise conjecture about the number of simple B -modules, $\ell(B)$, in terms of local information. If D is abelian, the conjecture reduces to

$$\ell(B) = \ell(b).$$

Broué's Conjecture

[B is a block of KG , defect group D , b its Brauer correspondent in $N_G(D)$.]

If D is abelian, Alperin's weight conjecture states that

$$\ell(B) = \ell(b);$$

is there a structural/geometric reason for B and b having the same number of simple modules?

Conjecture (Broué, 1990)

Let G be a finite group, and let B be a ℓ -block of G with abelian defect group D . Let b be the Brauer correspondent in $N_G(D)$. Then B and b are derived equivalent.

When Is Broué's Conjecture Known?

Broué's conjecture is known for quite a few groups:

- A_n, S_n (Chuang–Rouquier, Marcus);
- $GL_n(q)$, $\ell \nmid q$ (Chuang–Rouquier);
- D cyclic, $C_2 \times C_2$ (Rouquier, Erdmann, Rickard);
- G finite, $\ell = 2$, B **principal**;
- G finite, $\ell = 3$, $|P| = 9$, B principal (Koshitani, Kunugi, Miyachi, Okuyama, Waki);
- $SL_2(q)$, $\ell \mid q$ (Chuang, Kessar, Okuyama);
- various low-rank Lie type groups $L(q)$ with $\ell \nmid q$.

The Principal Block

If B_1, \dots, B_r are the blocks of KG , then the simple KG -modules are exactly the union of the simple B_i -modules.

The block contributing the trivial module is called the **principal block**, and denoted by $B_0(KG)$. Its defect group is always the Sylow ℓ -subgroup P , so its Brauer correspondent is a block of $KN_G(P)$.

Theorem (Brauer's third main theorem)

The Brauer correspondent of $B_0(KG)$ is $B_0(KN_G(P))$.

Thus if we are considering principal blocks, we need to relate the principal block of KG with the principal block of $KN_G(P)$.

Principal Blocks Are Good

In representation theory, one standard method of proof is to reduce a conjecture to finite simple groups and then use the Classification of the Finite Simple Groups.

In general, there is no (known) reduction of Broué's conjecture to simple groups, but for principal blocks there is.

Theorem

Let G be a finite group, and suppose that P is abelian. Then there are normal subgroups $H \leq L$ such that

- $\ell \nmid |H|$,
- $\ell \nmid |G : L|$, and
- L/H is a direct product of simple groups and an abelian ℓ -group.

For **principal** blocks, we may assume that $H = 1$. A derived equivalence for L passes up to G . Thus if Broué's conjecture for principal blocks holds for all simple groups, it holds for all groups.

How Do You Find Derived Equivalences?

There are four main methods to prove that B and b are derived equivalent.

- 1 **Okuyama deformations**: using many steps, deform the **Green correspondents of the** simple modules for B into those for b . This works well for small groups.
- 2 **Rickard's Theorem**: randomly find complexes in the derived category of b related to the **Green correspondents of the** simple modules for B , and if they 'look' like simple modules (**i.e., Homs and Exts behave nicely**) then there is a derived equivalence $B \rightarrow b$.
- 3 **More structure**: if B and b are more closely related (say **Morita** or **Puig** equivalent) then they are derived equivalent. More generally, find another block B' for some other group, an equivalence $B \rightarrow B'$, and a (previously known) equivalence $B' \rightarrow b$.
- 4 **Perverse equivalence**: build a derived equivalence up step by step in an algorithmic way.

What is a Perverse Equivalence?

Let A and B be finite-dimensional algebras.

An equivalence $F : D^b(\text{mod-}A) \rightarrow D^b(\text{mod-}B)$ is **perverse** if there exist

- orderings on the simple modules $S_1, S_2, \dots, S_r, T_1, T_2, \dots, T_r$, and
- a function $p : \{1, \dots, r\} \rightarrow \mathbb{Z}$

such that, if \mathcal{A}_i denotes the **Serre subcategory** generated by S_1, \dots, S_i , then

- F induces equivalences $D^b(\mathcal{A}_i) \rightarrow D^b(\mathcal{B}_i)$, and
- $F[p(i)]$ induces an equivalence $\mathcal{A}_i/\mathcal{A}_{i-1} \rightarrow \mathcal{B}_i/\mathcal{B}_{i-1}$.

Note that $\text{mod-}B$ is determined, up to equivalence, by A , p , and the ordering of the S_j .

Benefits of a Perverse Equivalence

The perverse equivalence is 'better' than a general derived equivalence.

- Has an underlying geometric interpretation (for Lie-type groups).
- The p -function comes from Lusztig's a -function (so is known).
- There is an algorithm that gives us a perverse equivalence from $B_0(KN)$ to **some** algebra, so only need to check that the target is $B_0(KG)$. (This is simply checking that the Green correspondents are the last terms in the complexes.)

This algorithm is very useful!

An Example

Let $G = M_{11}$, $\ell = 3$.

p	Ord. Char.	S_1	S_3	S_7	S_2	S_4	S_6	S_5
0	1	1						
2	10		1					
3	10			1				
4	16	1	1		1			
5	11	1			1	1		
6	44			1	1	1	1	
7	55	1	1		1	1	1	1
	10							1
	16	1				1		1

The cohomology of the complexes gives the rows of the decomposition matrix.

Which Groups Have Perverse Equivalences?

- $\mathrm{PSL}_3(q)$, $\ell = 3 \mid (q - 1)$
- $\mathrm{PSL}_4(q)$, $\mathrm{PSL}_5(q)$, $\ell = 3 \mid (q + 1)$
- $\mathrm{PSU}_3(q)$, $\ell = 3 \mid (q + 1)$
- $\mathrm{PSU}_4(q)$, $\mathrm{PSU}_5(q)$, $\ell = 3 \mid (q - 1)$
- $\mathrm{PSp}_4(q)$, $\ell = 3 \mid (q - 1), (q + 1)$
- (almost) $\mathrm{PSp}_8(q)$, $\ell = 5 \mid (q^2 + 1)$, $P = C_5 \times C_5$
- (almost) $\Omega_8^+(q)$, $\ell = 5 \mid (q^2 + 1)$, $P = C_5 \times C_5$
- $G_2(q)$, $\ell = 5 \mid (q + 1)$, $P = C_5 \times C_5$
- S_6 , A_7 , A_8 , $\ell = 3$ (A_6 does not)
- M_{11} , $M_{22.2}$, M_{23} , HS , $\ell = 3$ (M_{22} does not)
- $\mathrm{SL}_2(8)$, $\ell = 2$ in two steps

An Example: $\mathrm{PSL}_3(q)$

p	Ord. Char	S_1	S_5	S_2	S_3	S_4
0	1	1				
2	$q(q+1)$	1	1			
3	$(q+1)(q^2+q+1)/3$	1	1	1		
3	$(q+1)(q^2+q+1)/3$	1	1		1	
3	$(q+1)(q^2+q+1)/3$	1	1			1

		H^{-3}	H^{-2}	H^{-1}	Total
X_5 :	$0 \rightarrow \mathcal{P}(5) \rightarrow \mathcal{P}(234) \rightarrow C_5 \rightarrow 0.$		$1/5$	11	$5 - 1$
X_2 :	$0 \rightarrow \mathcal{P}(2) \rightarrow \mathcal{P}(34) \rightarrow \mathcal{P}(5) \rightarrow C_2 \rightarrow 0.$	$1/5/2$	1		$2 - 5$
X_3 :	$0 \rightarrow \mathcal{P}(3) \rightarrow \mathcal{P}(24) \rightarrow \mathcal{P}(5) \rightarrow C_3 \rightarrow 0.$	$1/5/3$	1		$3 - 5$
X_4 :	$0 \rightarrow \mathcal{P}(4) \rightarrow \mathcal{P}(23) \rightarrow \mathcal{P}(5) \rightarrow C_4 \rightarrow 0.$	$1/5/4$	1		$4 - 5$

An Example: $\mathrm{PSp}_4(q)$

p	Ord. Char	S_1	S_5	S_2	S_3	S_4
0	1	1				
3	$q(q-1)^2/2$		1			
3	$q(q^2+1)/2$	1		1		
3	$q(q^2+1)/2$	1			1	
4	q^4	1	1	1	1	1

$$X_5 : 0 \rightarrow \mathcal{P}(5) \rightarrow \mathcal{P}(234) \rightarrow M_{4,1} \oplus M_{4,2} \rightarrow C_5 \rightarrow 0.$$

$$X_2 : 0 \rightarrow \mathcal{P}(2) \rightarrow \mathcal{P}(5) \rightarrow \mathcal{P}(3) \oplus M_{1,2} \rightarrow C_2 \rightarrow 0.$$

$$X_3 : 0 \rightarrow \mathcal{P}(3) \rightarrow \mathcal{P}(5) \rightarrow \mathcal{P}(2) \oplus M_{1,1} \rightarrow C_3 \rightarrow 0.$$

$$X_4 : 0 \rightarrow \mathcal{P}(4) \rightarrow \mathcal{P}(4) \rightarrow \mathcal{P}(23) \rightarrow \mathcal{P}(5) \rightarrow C_4 \rightarrow 0.$$

An Example: $\mathrm{PSL}_4(q)$

p	Ord. Char.	S_1	S_2	S_5	S_3	S_4
0	1	1				
3	$q(q^2 + q + 1)$	1	1			
4	$q^2(q^2 + 1)$		1	1		
5	$q^3(q^2 + q + 1)$	1	1	1	1	
6	q^6	1			1	1

$$X_2 : \quad 0 \rightarrow \mathcal{P}(2) \rightarrow \mathcal{P}(5) \rightarrow \mathcal{P}(3) \oplus M_{1,2} \rightarrow C_2 \rightarrow 0.$$

$$X_5 : \quad 0 \rightarrow \mathcal{P}(5) \rightarrow \mathcal{P}(345) \rightarrow \mathcal{P}(234) \oplus M_{4,1} \rightarrow M_{4,1} \oplus M_{4,2} \rightarrow C_5 \rightarrow 0.$$

$$X_3 : \quad 0 \rightarrow \mathcal{P}(3) \rightarrow \mathcal{P}(34) \rightarrow \mathcal{P}(45) \rightarrow \mathcal{P}(5) \oplus M_{1,1} \rightarrow M_{1,1} \oplus M_{1,2} \rightarrow C_3 \rightarrow 0.$$

$$X_4 : \quad 0 \rightarrow \mathcal{P}(4) \rightarrow \mathcal{P}(4) \rightarrow \mathcal{P}(3) \rightarrow \mathcal{P}(3) \rightarrow \mathcal{P}(4) \rightarrow M_{4,2} \rightarrow C_4 \rightarrow 0.$$

Some Remarks

- Since $p(-)$, the ordering and the first category determine the perverse equivalence, it is a very compact way of defining a (type of) derived equivalence.
- Computationally, this reduces finding a derived equivalence to finding the Green correspondents of the simple modules for G , a much simpler task.
- For groups of Lie type, it seems as though the complexes above do not really depend on ℓ , and only that $\ell \mid \Phi_d(q)$. It might be possible to use these perverse equivalences to prove real results in this direction.