

Rainbow Matchings of Size $\delta(G)$ in Properly Edge-Colored Graphs

Jennifer Diemunsch^{1,4} Michael Ferrara^{1,5} Allan Lo^{2,6}
Casey Moffatt¹ Florian Pfender³ Paul S. Wenger¹

Submitted: Dec 21, 2011; Accepted: Jun 16, 2012; Published: Jun 28, 2012

Abstract

A *rainbow matching* in an edge-colored graph is a matching in which all the edges have distinct colors. Wang asked if there is a function $f(\delta)$ such that a properly edge-colored graph G with minimum degree δ and order at least $f(\delta)$ must have a rainbow matching of size δ . We answer this question in the affirmative; an extremal approach yields that $f(\delta) = 98\delta/23 < 4.27\delta$ suffices. Furthermore, we give an $O(\delta(G)|V(G)|^2)$ -time algorithm that generates such a matching in a properly edge-colored graph of order at least 6.5δ .

Keywords: Rainbow matching, properly edge-colored graphs

1 Introduction

All graphs under consideration in this paper are simple, and we let $\delta(G)$ and $\Delta(G)$ denote the minimum and maximum degree of a graph G , respectively. In this paper, we consider edge-colored graphs and let $c(uv)$ denote the color of the edge uv . An edge coloring of a graph is *proper* if the colors on edges sharing an endpoint are distinct. An edge-colored graph is *rainbow* if all edges have distinct colors. Rainbow matchings are of particular interest given their connection to transversals of Latin squares: each Latin square can be converted to a properly edge-colored complete bipartite graph, and a transversal of the Latin square is a rainbow perfect matching in the graph. Ryser's conjecture [9] that every Latin square of odd order has a transversal can be seen as the beginning of the study of rainbow

¹Dept. of Mathematical and Statistical Sciences, Univ. of Colorado Denver, Denver, CO; email addresses: jennifer.diemunsch@ucdenver.edu, michael.ferrara@ucdenver.edu, casey.moffatt@ucdenver.edu, paul.wenger@ucdenver.edu.

²School of Mathematics, University of Birmingham, Birmingham B15 2TT, UK, s.a.lo@bham.ac.uk

³Institut für Mathematik, Univ. Rostock, Rostock, Germany; Florian.Pfender@uni-rostock.de.

⁴Research supported in part by UCD GK12 Transforming Experiences Project, NSF award # 0742434.

⁵Research supported in part by Simons Foundation Collaboration Grant # 206692.

⁶Research supported in part by European Research Council Grant # 258345.

matchings. Stein [10] later conjectured that every Latin square of order n has a transversal of size $n - 1$; equivalently every proper edge-coloring $K_{n,n}$ has a rainbow matching of size $n - 1$. The connection between Latin transversals and rainbow matchings in $K_{n,n}$ has inspired additional interest in the study of rainbow matchings in triangle-free graphs. A thorough survey of rainbow matching and other rainbow subgraphs in edge-colored graphs appears in [6].

Several results have been attained for rainbow matchings in arbitrarily edge-colored graphs. The *color degree* of a vertex v in an edge-colored graph G , written $\hat{d}(v)$, is the number of distinct colors on edges incident to v . We let $\hat{\delta}(G)$ denote the minimum color degree among the vertices in G . Wang and Li [12] proved that every edge-colored graph G contains a rainbow matching of size at least $\left\lceil \frac{5\hat{\delta}(G)-3}{12} \right\rceil$, and conjectured that a rainbow matching of size $\lceil \hat{\delta}(G)/2 \rceil$ exists if $\hat{\delta}(G) \geq 4$. LeSaulnier et al. [8] then proved that every edge-colored graph G contains a rainbow matching of size $\lfloor \hat{\delta}(G)/2 \rfloor$. Finally, Kostochka and Yancey [5] proved the conjecture of Wang and Li in full, and also that triangle-free graphs have rainbow matchings of size $\lceil 2\hat{\delta}(G)/3 \rceil$.

Since the edge-colored graphs generated by Latin squares are properly edge-colored, it is of interest to consider rainbow matchings in properly edge-colored graphs. In this direction, LeSaulnier et al. proved that a properly edge-colored graph G satisfying $|V(G)| \neq \delta(G) + 2$ that is not K_4 has a rainbow matching of size $\lceil \delta(G)/2 \rceil$. Wang then asked if there is a function f such that a properly edge-colored graph G with minimum degree δ and order at least $f(\delta)$ must contain a rainbow matching of size δ [11]. As a first step towards answering this question, Wang showed that a graph G with order at least $8\delta/5$ has a rainbow matching of size $\lfloor 3\delta(G)/5 \rfloor$.

Since there are $n \times n$ Latin squares with no transversals when n is even (see [1, 14]), for such a function f it is clear that $f(\delta) > 2\delta$ when δ is even. Furthermore, since maximum matchings in $K_{\delta,n-\delta}$ have only δ edges (provided $n \geq 2\delta$), there is no function for the order of G depending on $\delta(G)$ that can guarantee a rainbow matching of size greater than $\delta(G)$.

In this paper we answer Wang's question from [11] in the affirmative, proving that a linear number of vertices in terms of the minimum degree suffices.

Theorem 1. *If G is a properly edge-colored graph satisfying $|V(G)| \geq 98\delta(G)/23$, then G contains a rainbow matching of size $\delta(G)$.*

Independently, Wang, Zhang, and Liu [13] also answering Wang's question in the affirmative, proving that a properly edge-colored graph G with at least $(\delta(G)^2 + 4\delta(G) - 4)/4$ vertices has a rainbow matching of size $\delta(G)$.

The proof of Theorem 1 utilizes extremal techniques akin to those that appear in [5, 8, 11] and [12]. We also implement a greedy algorithmic approach to demonstrate that it is possible to efficiently construct a rainbow matching of size δ in a properly edge-colored graph with minimum degree δ having order at least 6.5δ . To our knowledge, an algorithmic approach of this type has not been previously employed in the study of rainbow matchings.

Theorem 2. *If G is a properly edge-colored graph with minimum degree δ satisfying $|V(G)| > \frac{13}{2}\delta - \frac{23}{2} + \frac{41}{8\delta}$, then there is an $O(\delta(G)|V(G)|^2)$ -time algorithm that produces a rainbow matching of size δ in G .*

As a contrast, Itai, Rodeh, and Tanimota [4] proved that determining if an edge-colored graph G has a rainbow matching of size k is NP-complete, even if G is bipartite. More recently, Le and Pfender [7] have shown that the problem of determining the maximum size of a rainbow matching in a properly edge-colored graph is NP-hard, even when restricted to properly edge-colored paths.

2 Proof of Theorem 1

Let G be a properly edge-colored n -vertex graph with minimum degree δ and $n \geq 98\delta/23$. The theorem holds easily if $\delta \leq 2$, so we may assume that $\delta \geq 3$. By way of contradiction, let G be a counterexample with δ minimized; thus G does not contain a rainbow matching of size δ . Further, we may assume that $|E(G)|$ is minimized, so in particular the vertices of degree greater than δ form an independent set, as otherwise we could delete an edge without lowering the minimum degree. We break the proof into a series of simple claims.

Let $\Delta(G) = d_1 \geq d_2 \geq \dots \geq d_n = \delta$ with $d(v_i) = d_i$ be the degree sequence of G .

Lemma 3. *For $1 \leq k \leq 2\delta/3$, there exists an $i \leq k$ such that $d_i \leq 3\delta - k - 2i$.*

Proof. Suppose that for some $k \leq 2\delta/3$, $d_i \geq 3\delta + 1 - k - 2i$ for all $1 \leq i \leq k$. It follows that $d_i > \delta$ for $i \leq k$, and therefore $\{v_1, \dots, v_k\}$ is an independent set. Delete the vertices v_1, v_2, \dots, v_k from G , and note that $\delta(G \setminus \{v_1, \dots, v_k\}) \geq \delta - k$. By the minimality of G , there exists a rainbow matching M_k of size $\delta - k$ in $G \setminus \{v_1, \dots, v_k\}$.

The vertex v_k has at most $2(\delta - k)$ neighbors in M_k , and is incident to at most $\delta - k$ edges colored with colors occurring in M_k . Thus, v_k has a neighbor $w_k \notin V(M_k)$ such that the color of $v_k w_k$ does not occur in M_k , and we can extend M_k by adding the edge $v_k w_k$; call the resulting rainbow matching M_{k-1} . Note that $w_k \neq v_i$ for $i \leq k$ as $\{v_1, \dots, v_k\}$ is an independent set.

The j th iteration of this process produces a rainbow matching M_{k-j} of size $\delta - k + j$ that contains $\{v_k, \dots, v_{k-j+1}\}$. Hence v_{k-j} has at most $2(\delta - k) + j$ neighbors in M_{k-j} and is incident to at most $\delta - k + j$ edges colored with colors occurring in M_{k-j} . Thus there is a vertex $w_{k-j} \in N(v_{k-j})$ such that the edge $v_{k-j} w_{k-j}$ extends M_{k-j} to a $(\delta - k + j + 1)$ -edge rainbow matching $M_{k-(j+1)}$. Continuing in this fashion extends the matching M_k by k edges, yielding a rainbow matching of size δ , a contradiction finishing the proof. \square

As a corollary of Lemma 3, we obtain the following.

Lemma 4. *For $1 \leq k \leq 2\delta/3$, we have $\sum_{i=1}^k d_i \leq k(3\delta - 2 - k)$, with equality only if $d_1 = d_k = 3\delta - 2 - k$.*

Proof. We proceed by induction on k . For $k = 1$, the statement follows from Lemma 3. Now let $k > 1$ and let $i \leq k$ such that $d_i \leq 3\delta - k - 2i$. By induction,

$$\sum_{j=1}^{i-1} d_j \leq (i-1)(3\delta - 1 - i), \text{ and}$$

$$\sum_{j=i}^k d_j \leq (k-i+1)d_i \leq (k-i+1)(3\delta - k - 2i).$$

Thus,

$$\sum_{j=1}^k d_j \leq (i-1)(3\delta - 1 - i) + (k-i+1)(3\delta - k - 2i) = 3k\delta - k^2 - k + 1 - i(k+2-i) \leq k(3\delta - 2 - k)$$

and equality holds only if $i = 1$ and $d_1 = d_k = 3\delta - 2 - k$. \square

Let C be a maximum color class in G and let $|C| = a$. By the minimality of G , there exists a rainbow matching $M = \{x_i y_i : 1 \leq i \leq \delta - 1\}$ of size $\delta - 1$ in $G - C$. Without loss of generality, we may assume that $c(x_i y_i) = i$ for $1 \leq i \leq \delta - 1$ and the edges in C have color δ . Let $W = V(G) \setminus V(M)$; observe that $|W| = n - 2(\delta - 1)$. If there is an edge e in $G[W]$ with $c(e) \notin \{1, \dots, \delta - 1\}$ then we can add e to M to obtain a rainbow matching of size δ . Thus we may assume that every edge whose color is not in $\{1, \dots, \delta - 1\}$ has an endpoint in $V(M)$. An edge uv is *good* if its color is not in $\{1, \dots, \delta - 1\}$ and one of its endpoints is in W . A vertex $v \in V(M)$ is *good* if v is incident with at least seven good edges.

Claim 5. *For $i \in \{1, \dots, \delta - 1\}$, if x_i is incident with at least three good edges, then no good edge is incident with y_i , and vice versa.*

Proof. Suppose that $y_i u$ is a good edge. If x_i is incident with at least three good edges, then x has a neighbor w such that vw is a good edge, $w \neq u$, and $c(x_i w) \neq c(y_i u)$. Thus $(M \cup \{x_i w, y_i u\}) \setminus \{x_i y_i\}$ is a rainbow matching of size δ , a contradiction. \square

By Claim 5, we may assume without loss of generality that $\{x_1, \dots, x_r\}$ is the set of good vertices for some $r \geq 0$. Let $W' = W \cup \{y_1, \dots, y_r\}$.

Claim 6. *No edge uv in $G[W']$ has color in $\{1, \dots, r\}$.*

Proof. By way of contradiction, assume that there is an edge uv in $G[W']$ such that $c(uv) \in \{1, \dots, r\}$. Let M' be the subset of M consisting of the edge with color $c(uv)$ and any edges with an endpoint in $\{u, v\}$. There are at most three such edges (the edge with color $c(uv)$ and possibly one for each endpoint); without loss of generality, let $M' = \{x_1 y_1, \dots, x_t y_t\}$ (here $1 \leq t \leq 3$). Note that x_j is a good vertex for $1 \leq j \leq t$. Thus there are distinct vertices w_1, \dots, w_t such that $x_j w_j$ is a good edge for $1 \leq j \leq t$ and the colors on the edges $uv, x_1 w_1, \dots, x_t w_t$ are distinct. Thus $(M \cup \{uv, x_1 w_1, \dots, x_t w_t\}) \setminus \{x_1 y_1, \dots, x_t y_t\}$ is a rainbow matching of size δ , a contradiction. \square

An edge uv is *nice* if its color is not in $\{r+1, \dots, \delta-1\}$ and one of its endpoints is in W' . Note that every good edge is nice. Recall that every good edge has an endpoint in $V(M)$. By Claim 5 and Claim 6, no nice edge lies in $G[W']$. Hence, every nice edge joins vertices in W' and $V(G)\setminus W'$. A vertex $v \in V(M)\setminus\{x_1, \dots, x_r, y_1, \dots, y_r\}$ is *nice* if v is incident with at least seven nice edges. Note that if there is no good vertex (i.e. $r = 0$), then the definitions of good and nice vertices are the same and so there is also no nice vertex. Next, we prove analogues of Claim 5 and Claim 6 for nice vertices and edges.

Claim 7. *For $i \in \{r+1, \dots, \delta-1\}$, if x_i is incident with at least three nice edges, then no nice edge is incident with y_i , and vice versa.*

Proof. Suppose $y_i u$ is a nice edge for some $i \in \{r+1, \dots, \delta-1\}$. If x_i is incident to at least three nice edges, then x_i has a neighbor v such that $x_i v$ is a nice edge, $v \neq u$, and $c(x_i v) \neq c(y_i u)$. Let M' be the subset of M consisting of edges with an endpoint in $\{u, v\}$ or a color in $\{c(x_i v), c(y_i u)\}$. There are at most four such edges (possibly one with each endpoint and one with each color); without loss of generality, let $M' = \{x_1 y_1, \dots, x_t y_t\}$ (here $0 \leq t \leq 4$). Note that x_j is a good vertex for $1 \leq j \leq t$. Thus there are distinct vertices w_1, \dots, w_t such that $x_j w_j$ is a good edge for $1 \leq j \leq t$ and the colors on the edges $x_i v, y_i u, x_1 w_1, \dots, x_t w_t$ are distinct. Thus $(M \cup \{x_i v, y_i u, x_1 w_1, \dots, x_t w_t\}) \setminus \{x_i y_i, x_1 y_1, \dots, x_t y_t\}$ is a rainbow matching of size δ , a contradiction. \square

By Claim 7, we may assume that $\{x_{r+1}, x_{r+2}, \dots, x_{r+s}\}$ is the set of nice vertices for some $s \geq 0$.

Claim 8. *No edge uv in $G[W']$ has color in $\{1, \dots, r+s\}$.*

Proof. By Claim 6, the claim holds if $s = 0$. Assume that $s \geq 1$, and consequently $r \geq 1$. Without loss of generality, suppose that there is an edge uv in $G[W']$ with $c(uv) = r+1$. Because x_{r+1} is nice, it has a neighbor v' in W' such that $x_{r+1} v'$ is a nice edge and $v' \neq u, v$. Let M' be the subset of M consisting of those edges an endpoint in $\{u, v, v'\}$ or color $c(x_{r+1} v')$. Again there are at most four edges in M' and we let $M' = \{x_1 y_1, \dots, x_t y_t\}$. Defining w_1, \dots, w_t as before, $(M \cup \{uv, x_{r+1} v', x_1 w_1, \dots, x_t w_t\}) \setminus \{x_{r+1} y_{r+1}, x_1 y_1, \dots, x_t y_t\}$ is a rainbow matching of size δ , a contradiction. \square

Next, we count the number of nice edges in G .

Claim 9. *There are at most $\max\{(3\delta - 8 - r + s)r + 6(\delta - 1), (7\delta/3 - 7 + s)r + 6(\delta - 1)\}$ nice edges in G .*

Proof. Recall that $V(G) \setminus W' = \{x_1, \dots, x_{\delta-1}, y_{r+1}, \dots, y_{\delta-1}\}$ and every nice edge joins vertices from W' and $V(G) \setminus W'$. If $r \leq 2\delta/3$, then the set of good vertices is incident to at most $r(3\delta - 2 - r)$ nice edges by Lemma 4. Similarly, if $r \geq 2\delta/3$, then the set of good vertices is incident to at most $r(3\delta - 2 - \lfloor 2\delta/3 \rfloor) \leq r(7\delta/3 - 1)$ nice edges. For $i \in \{r+1, \dots, r+s\}$, Claim 7 implies that x_i is incident to at most $r+6$ nice edges and y_i is incident to none.

For $i \in \{r + s + 1, \dots, \delta - 1\}$, by Claim 7 there are at most six nice edges with an endpoint in $\{x_i, y_i\}$. Therefore, the number of nice edges is at most

$$(3\delta - 2 - r)r + (r + 6)s + 6(\delta - 1 - r - s) = (3\delta - 8 - r + s)r + 6(\delta - 1) \text{ if } r \leq 2\delta/3,$$

and

$$(7\delta/3 - 1)r + (r + 6)s + 6(\delta - 1 - r - s) = (7\delta/3 - 7 + s)r + 6(\delta - 1) \text{ if } r \geq 2\delta/3. \quad \square$$

Recall that C is the color class with color δ , $|C| = a$, and C is a maximum size color class. Therefore there are at least $2(a - \delta + 1)$ vertices in W incident to an edge in C . Since every edge in C has an endpoint in $V(M)$ it follows that there are at least $2(a - \delta + 1)$ vertices in $V(M)$ joined to W by edges in C . Without loss of generality, let $\{r + s + 1, \dots, r + s + t\}$ be the set of indices $i \in \{r + s + 1, \dots, \delta - 1\}$ such that x_i or y_i is incident to an edge in C . By Claim 5 and Claim 7, we have

$$t \geq a - \delta + 1 - \frac{r + s}{2} \text{ and } r + s + t \leq \delta - 1. \quad (1)$$

Claim 10. For $i \in \{r + s + 1, \dots, r + s + t\}$, there is at most one edge of color i in $G[W]$.

Proof. Suppose uv and $u'v'$ are edges of color i in $G[W]$ for some $i \in \{r + s + 1, \dots, r + s + t\}$. Without loss of generality, we may assume that there exists $w \in W$ such that $c(x_i w) = \delta$ and $w \neq u, v$. Hence, $(M \cup \{uv, x_i w\}) \setminus \{x_i y_i\}$ is a rainbow matching of size δ , a contradiction. \square

Now we count the number of nice edges from W' to $V(G) \setminus W'$. Recall that each color class in G contains at most a edges. By Claim 8, there is no edge in $G[W']$ of color $i \in \{r + 1, \dots, r + s\}$. Thus, for $i \in \{r + 1, \dots, r + s\}$ there are at most $a - 1$ vertices in W' that are incident with an edge of color i . Since every color class has size at most a , for $i \in \{r + s + 1, \dots, \delta - 1\}$ there are at most $2(a - 1)$ vertices in W' that are incident with an edge of color i . Furthermore for $i \in \{r + s + 1, \dots, r + s + t\}$, Claim 10 implies that there are at most a vertices in W that are incident with an edge of color i . Since $W' \setminus W = \{y_1, \dots, y_r\}$, it follows that for $i \in \{r + s + 1, \dots, r + s + t\}$ there are at most $\min\{a + r, 2(a - 1)\}$ vertices in W' that are incident with an edge of color i . It then follows, using the fact that $|W'| = |W| + r = n - 2(\delta - 1) + r$ and (1), that the number of nice edges from W' to $V(G) \setminus W'$ is at least

$$\begin{aligned} & \delta|W'| - (a - 1)(2\delta - 2 - 2r - s - 2t) - \min\{a + r, 2(a - 1)\}t \\ &= \delta n - 2\delta(\delta - 1) + \delta r - (a - 1)(2\delta - 2 - 2r - s) + \max\{a - r - 2, 0\}t. \end{aligned}$$

We first consider the case when $r \leq 2\delta/3$. Applying the upper bound of $(3\delta - 8 - r + s)r + 6(\delta - 1)$ nice edges from Claim 9, we obtain

$$\delta n \leq (2\delta - 8 - r + s)r + 2(\delta + 3)(\delta - 1) + (a - 1)(2\delta - 2 - 2r - s) - \max\{a - r - 2, 0\}t. \quad (2)$$

To finish the proof we bound the right hand side of (2) to obtain a contradiction. Note that the coefficient of t is nonpositive. Thus, the right hand side of (2) is maximized when t is minimized. By (1), $t \geq \max\{a - \delta + 1 - (r + s)/2, 0\}$.

If $a \leq \delta - 1 + (r + s)/2$, then we let $t = 0$. The coefficient of a is nonnegative, and thus (2) is maximized when a is maximized; hence we assume $a = \delta - 1 + (r + s)/2$. Substituting for a yields a negative quadratic in s that is maximized when $s = 1 - r/2$. Since s is a nonnegative integer and $s = 0$ if $r = 0$, (2) is maximized at $s = 0$, which yields

$$\delta n \leq 2(2\delta + 1)(\delta - 1) + (\delta - 5 - 2r)r.$$

This is maximized when $r = (\delta - 5)/4$, yielding $n \leq 33\delta/8 - 13/4 + 9/(8\delta)$, a contradiction.

If $a > \delta - 1 + r/2$, we let $t = a - \delta + 1 - r/2$. Since $t > 0$, it follows that $r + s \leq \delta - 2$. Thus $a - r - 2 > \delta - 3 - (r - s)/2 \geq \delta - 3 - (\delta - 2)/2 = \delta/2 - 2$. As $\delta \geq 3$ and $a - r - 2$ is an integer, $\max\{a - r - 2, 0\} = a - r - 2$. Therefore the coefficient of s in (2) is nonpositive and we may assume that $s = 0$. Consequently, (2) becomes

$$\delta n \leq (3\delta - 1 - r/2 - a)a + (\delta - 6 - 3r/2)r + 2(\delta^2 - 1).$$

The right hand side is maximized when $a = 3\delta/2 - 1/2 - r/4$, so

$$\delta n \leq \frac{1}{16}(-28 + 68\delta^2 - 24\delta + 4\delta r - 92r - 23r^2),$$

which is maximized when $r = 2\delta/23 - 2$. This yields

$$n \leq 98\delta/23 - 2 + 4/\delta,$$

a contradiction.

To complete the proof of the theorem, we are left with the case $r \geq 2\delta/3$. Similarly to (2), since we have at most $(7\delta/3 - 7 + s)r + 6(\delta - 1)$ nice edges in G by Claim 9, we have

$$\delta n \leq (4\delta/3 - 7 + s)r + 2(\delta + 3)(\delta - 1) + (a - 1)(2\delta - 2 - 2r - s) - \max\{a - r - 2, 0\}t. \quad (3)$$

Again, the right hand side of (3) is maximized when t is minimized.

If $a \leq \delta - 1 + (r + s)/2$, then (3) is maximized when $t = 0$ and $a = \delta - 1 + r/2$. Again we may assume that $s = 0$, yielding

$$\delta n \leq -2 + 4\delta^2 - 2\delta + r(\delta/3 - 4 - r).$$

This is maximized when $r = \delta/6 - 2$, yielding $n \leq 145\delta/36 - 8/3 + 6/\delta$, a contradiction.

If $a > \delta - 1 + (r + s)/2$, we let $t = a - \delta + 1 - (r + s)/2$ and again we may assume that $s = 0$. Then, (3) becomes

$$\delta n \leq \frac{1}{6}(-6a^2 + 3a(6\delta - 2) + 12\delta^2 - (3a + 30 + 3r - 2\delta)r - 12),$$

which is maximized when r is minimized. Since we have assumed that $r \geq 2\delta/3$, we have $r = 2\delta/3$ and we are back in the case $r \leq 2\delta/3$, finishing the proof of the theorem. \square

3 Proof of Theorem 2

We proceed by induction on $\delta(G)$. The result is trivial if $\delta(G) = 1$. We assume that G is a graph with minimum degree $\delta > 1$ and order greater than $\frac{13}{2}\delta - \frac{23}{2} + \frac{41}{8\delta}$.

Lemma 11. *If G has a color class containing at least $2\delta - 1$ edges, then G has a rainbow matching of size δ .*

Proof. Let C be a color class with at least $2\delta - 1$ edges. By induction, there is a rainbow matching M of size $\delta - 1$ in $G - C$. There are $2\delta - 2$ vertices covered by the edges in M , thus one of the edges in C has no endpoint covered by M , and the matching can be extended. \square

It is also useful to note that we also have the following, which is identical to Lemma 3 when $k = 1$.

Lemma 12. *If G satisfies $\Delta(G) > 3\delta - 3$, then G has a rainbow matching of size δ .*

We begin by preprocessing the graph so that each edge is incident to at least one vertex with degree δ . To achieve this, arbitrarily order the edges in G and process them in order. If both endpoints of an edge have degree greater than δ when it is processed, delete that edge. In the resulting graph, every edge is incident to a vertex with degree δ . Furthermore, by Lemma 12 we may assume that $\Delta(G) \leq 3\delta - 3$; thus the degree sum of the endpoints of any edge is bounded above by $4\delta - 3$. After preprocessing, we begin the greedy algorithm.

In the i th step of the algorithm, a smallest color class is chosen (without loss of generality, color i), and then an edge e_i of color i is chosen such that the degree sum of the endpoints is minimized. All the remaining edges of color i and all edges incident with the endpoints of e_i are deleted. The algorithm terminates when there are no edges in the graph.

We assume that the algorithm fails to produce a matching of size δ in G ; suppose that the rainbow matching M generated by the algorithm has size k . We let R denote the set of vertices that are not covered by M .

Let c_i denote the size of the smallest color class at step i . Since at most two edges of color $i + 1$ are deleted in step i (one at each endpoint of e_i), we observe that $c_{i+1} + 2 \geq c_i$. Otherwise, at step i color class $i + 1$ has fewer edges. Let step h be the last step in the algorithm in which a color class that does not appear in M is completely removed from G . It then follows that $c_h \leq 2$, and in general $c_i \leq 2(h - i + 1)$ for $i \in [h]$. Let f_i denote the number of edges of color i deleted in step i with both endpoints in R . Since $f_i < c_i$, we have $f_i \leq 2(h - i) + 1$ for $i \in [h]$. Note that after step h , there are exactly $k - h$ colors remaining in G . By Lemma 11, color classes contain at most $2\delta - 2$ edges, and therefore the last $k - h$ steps remove at most $(k - h)(2\delta - 2)$ edges. Furthermore, for $i > h$, the degree sum of the endpoints of e_i is at most $2(\delta - 1)$.

For $i \in [h]$, let x_i and y_i be the endpoints of e_i , and let $d_i(v)$ denote the degree of a vertex v at the beginning of step i . Let $\mu_i = \max\{0, d_i(x_i) + d_i(y_i) - 2\delta\}$; note that $2\delta \leq 2\delta + \mu_i \leq 4\delta - 3$. Thus, at step i , at most $2\delta + \mu_i + f_i - 1$ edges are removed from the

graph. Since the algorithm removes every edge from the graph, we conclude that

$$|E(G)| \leq (k-h)(2\delta-2) + \sum_{i=1}^h (2\delta + \mu_i + f_i - 1). \quad (4)$$

We now compute a lower bound for the number of edges in G . Since the degree sum of the endpoints of e_i in G is at least $2\delta + \mu_i$, we immediately obtain the following inequality:

$$\frac{n\delta + \sum_{i \in [h]} \mu_i}{2} \leq |E(G)|.$$

If $f_i > 0$ and $\mu_i > 0$, then there is an edge with color i having both endpoints in R . Since this edge was not chosen in step i by the algorithm, the degree sum of its endpoints is at least $2\delta + \mu_i$, and one of its endpoints has degree at least $\delta + \mu_i$. For each value of i satisfying $f_i > 0$, we wish to choose a representative vertex in R with degree at least $\delta + \mu_i$. Since there are f_i edges with color i having both endpoints in R , there are f_i possible representatives for color i . Since a vertex in R with high degree may be the representative for multiple colors, we wish to select the largest system of distinct representatives.

Suppose that the largest system of distinct representatives has size t , and let T be the set of indices of the colors that have representatives. For each color $i \in T$ there is a distinct vertex in R with degree at least $\delta + \mu_i$. Thus we may increase the edge count of G as follows:

$$\frac{n\delta + \sum_{i \in [h]} \mu_i + \sum_{i \in T} \mu_i}{2} \leq |E(G)|. \quad (5)$$

We let $\{f_i^\downarrow\}$ denote the sequence $\{f_i\}_{i \in [h]}$ sorted in nonincreasing order. Since $f_i \leq 2(h-i)+1$, we conclude that $f_i^\downarrow \leq 2(h-i)+1$. Because there is no system of distinct representatives of size $t+1$, the sequence $\{f_i^\downarrow\}$ cannot majorize the sequence $\{t+1, t, t-1, \dots, 1\}$. Hence there is a smallest value $p \in [t+1]$ such that $f_p^\downarrow \leq t+1-p$. Therefore, the maximum value of $\sum_{i=1}^h f_i^\downarrow$ is bounded by the sum of the sequence $\{2h-1, 2h-3, \dots, 2(h-p)+3, t+1-p, \dots, t+1-p\}$. Summing we attain

$$\sum_{i \in [h]} f_i \leq (p-1)(2h-p+1) + (h-p+1)(t+1-p).$$

Over p , this value is maximized when $p = t+1$, yielding $\sum_{i \in [h]} f_i \leq t(2h-t)$. Since $h \leq \delta-1$, we then have $\sum_{i \in [h]} f_i \leq 2(\delta-1)t - t^2$.

We now combine bounds (4) and (5):

$$\frac{n\delta + \sum_{i \in [h]} \mu_i + \sum_{i \in T} \mu_i}{2} \leq (k-h)(2\delta-2) + \sum_{i=1}^h (2\delta + \mu_i + f_i - 1).$$

Hence, since $k \leq \delta - 1$,

$$\begin{aligned} \frac{n\delta}{2} &\leq (2\delta - 1)(\delta - 1) + \frac{1}{2} \sum_{[h] \setminus T} \mu_i + \sum_{i \in [h]} f_i \\ &\leq (2\delta - 1)(\delta - 1) + (\delta - 1 - t)(\delta - 3/2) + 2(\delta - 1)t - t^2 \\ &\leq 3\delta^2 - \frac{11}{2}\delta + \frac{5}{2} + \left(\delta - \frac{1}{2}\right)t - t^2. \end{aligned}$$

This bound is maximized when $t = (\delta - \frac{1}{2})/2$. Thus

$$n \leq \frac{13}{2}\delta - \frac{23}{2} + \frac{41}{8\delta},$$

contradicting our choice for the order of G .

It remains to show that this proof provides the framework of a $O(\delta(G)|V(G)|^2)$ -time algorithm that generates a rainbow matching of size $\delta(G)$ in a properly edge-colored graph G of order at least $\frac{13}{2}\delta - \frac{23}{2} + \frac{41}{8\delta}$. Given such a G , we create a sequence of graphs $\{G_i\}$ as follows, letting $G = G_0$, $\delta = \delta(G)$, and $n = |V(G)|$. First, determine $\delta(G_i)$, $\Delta(G_i)$, and the maximum size of a color class in G_i ; this process takes $O(n^2)$ -time. If $\Delta(G_i) \leq 3\delta(G_i) - 3$ and the maximum color class has at most $2\delta(G_i) - 2$ edges, then terminate the sequence and set $G_i = G'$. If $\Delta(G_i) > 3\delta(G_i) - 3$, then delete a vertex v of maximum degree and then process the edges of $G_i - v$, iteratively deleting those with two endpoints of degree at least $\delta(G_i)$; the resulting graph is G_{i+1} . If $\Delta(G_i) \leq 3\delta(G_i) - 3$ but a maximum color class C has at least $2\delta(G_i) - 1$ edges, then delete C and then process the edges of $G_i - C$, iteratively deleting those with two endpoints of degree at least $\delta(G_i)$; the resulting graph is G_{i+1} . Note that $\delta(G_{i+1}) = \delta(G_i) - 1$. If this process generates G_δ , we set $G' = G_\delta$ and terminate. Generating the sequence $\{G_i\}$ consists of at most δ steps, each taking $O(n^2)$ -time.

Given that $G' = G_i$, the algorithm from the proof of Theorem 2 takes $O(\delta n^2)$ -time to generate a matching of size $\delta - i$ in G' . The preprocessing step and the process of determining a smallest color class and choosing an edge in that class whose endpoints have minimum degree sum both take $O(n^2)$ -time. This process is repeated at most δ times.

A matching of size $\delta - (i + 1)$ in G_{i+1} is easily extended in G_i to a matching of size $\delta - i$ using the vertex of maximum degree or maximum color class. The process of extending the matching takes $O(\delta)$ -time. Thus the total run-time of the algorithm generating the rainbow matching of size δ in G is $O(\delta n^2)$. \square

It is worth noting that the analysis of the greedy algorithm used in the proof of Theorem 2 could be improved. In particular, the bound $c_{i+1} \geq c_i - 2$ is sharp only if at step i there are an equal number of edges of color i and $i + 1$ and both endpoints of e_i are incident to edges with color $i + 1$. However, since one of the endpoints of e_i has degree at most δ , at most $\delta - 1$ color classes can lose two edges in step i . Since the maximum size of a color class in G is at most $2\delta - 2$, if G has order at least 6δ , then there are at least $3\delta/2$ color classes. Thus, for small values of i , the bound $c_i \leq 2(k - i + 1)$ can likely be improved. However, we doubt that such analysis of this algorithm can be improved to yield a bound on $|V(G)|$ better than 6δ .

References

- [1] R. A. Brualdi and H. J. Ryser, *Combinatorial Matrix Theory*, Cambridge University Press, Cambridge, UK, 1991.
- [2] R. Diestel, *Graph Theory*, Springer-Verlag, New York, 1997.
- [3] G. A. Dirac, Some theorems on abstract graphs, *Proc. London Math. Soc.* (3) **2** (1952), 69-81.
- [4] A. Itai, M. Rodeh, and S. Tanimoto, Some matching problems for bipartite graphs, *J. Assoc. Comput. Mach.* **25** (1978), no. 4, 517-525.
- [5] A. Kostochka and M. Yancey, Large rainbow matchings in edge-colored graphs, In Preparation.
- [6] M. Kano and X. Li, Monochromatic and heterochromatic subgraphs in edge-colored graphs – a survey, *Graphs Comb.* **24** (2008), 237–263.
- [7] V. B. Le and F. Pfender, Complexity results for rainbow matchings, In Preparation.
- [8] T. D. LeSaulnier, C. Stocker, P. S. Wenger, and D. B. West, Rainbow matching in edge-colored graphs, *Electron. J. Combin.* **17** (2010), #N26.
- [9] H. J. Ryser, Neuere Probleme der Kombinatorik, in “Vorträge über Kombinatorik Oberwolfach”, Mathematisches Forschungsinstitut Oberwolfach, July 1967, 24–29.
- [10] S. K. Stein, Transversals of Latin squares and their generalizations, *Pacific J. Math.* **59** (1975), no. 2, 567–575.
- [11] G. Wang, Rainbow matchings in properly edge colored graphs, *Electron. J. Combin.* **18** (2011), #P162.
- [12] G. Wang and H. Li, Heterochromatic matchings in edge-colored graphs, *Electron. J. Combin.* **15** (2008), #R138.
- [13] G. Wang, J. Zhang, and G. Liu, Existence of rainbow matchings in properly edge-colored graphs, *Front. Math. China* **7** (2012), no. 3, 543–550.
- [14] I. M. Wanless, Transversals in Latin squares: a survey, *Surveys in Combinatorics 2011*, London Math. Soc. Lecture Note Series **392** (2011), 403–437.